

## SIGNED GENERALIZED PETERSEN GRAPH AND ITS CHARACTERISTIC POLYNOMIAL

E. GHASEMIAN AND G. H. FATH-TABAR\*

ABSTRACT. Let  $G^\sigma$  be a signed graph, where  $G = (V, E)$  is the underlying simple graph and  $\sigma : E(G) \rightarrow \{\pm 1\}$  is the sign function on  $E(G)$ . In this paper, we obtain  $k$ -th signed spectral moments and  $k$ -th signed Laplacian spectral moments of a signed graph  $G^\sigma$ , together with coefficients of their signed characteristic polynomial and signed Laplacian characteristic polynomial are calculated.

### 1. INTRODUCTION

Let  $G = (V, E)$  be a simple graph and  $\sigma : E(G) \rightarrow \{\pm 1\}$  a mapping on the edge set of  $G$ . The graph  $G$  together with the sign function  $\sigma$  is called a *signed graph* and is denoted by  $G^\sigma$ . If  $\sigma(e) = +1$ , then the edge  $e$  is called *positive*, and if  $\sigma(e) = -1$ , then the edge  $e$  is called *negative*. A *walk of length  $k$*  in a graph  $G$  is a sequence  $v_1 e_1 \dots v_k e_k v_{k+1}$  with vertices  $v_1, \dots, v_{k+1}$  and edges  $e_1, \dots, e_k$  such that, we have  $v_i \neq v_{i+1}$ ,  $1 \leq i \leq k$  and  $e_i$  is an edge from  $v_i$  to  $v_{i+1}$ . In a signed graph  $G^\sigma$ , a walk is called *positive* (resp., *negative*) if the number of its negative edges is even (resp., odd). The number of positive (resp., negative) walks of length  $k$  from vertex  $v_i$  to vertex  $v_j$  is denoted by  $w_{ij}^+(k)$  (resp.,  $w_{ij}^-(k)$ ). A *cycle* with  $n \geq 3$  vertices is a simple graph whose vertices can be sorted as a sequence such that two vertices are adjacent if and only if they are subsequent members of the sequence. In a signed graph  $G^\sigma$ , a cycle is called *balanced* or *positive* (resp., *unbalanced* or *negative*) if the number of its negative edges is even (resp., odd). A signed graph

---

MSC(2010): Primary: 05C50; Secondary: 15A18.

Keywords: Signed graph, signed generalized Petersen graph, adjacency matrix.

Received: 04 March 2017, Accepted: 01 December 2017.

\*Corresponding author.

is called *balanced* if all its cycles are balanced; otherwise, it is called *unbalanced* [1].

The *adjacency matrix* of a graph  $G$ , denoted by  $A(G) = [a_{ij}]$ , is an  $n \times n$  matrix where  $a_{ij} = 1$  if  $v_i v_j$  is an edge of the graph and  $a_{ij} = 0$  otherwise. The *adjacency matrix of a signed graph*, denoted by  $A(G^\sigma) = [a_{ij}^\sigma]$ , is an  $n \times n$  matrix where  $a_{ij}^\sigma = \sigma(ij)a_{ij}$  if  $v_i v_j$  is an edge of the graph, and  $a_{ij}^\sigma = 0$  otherwise. Thus, if  $e = ij$  is an edge and  $\sigma(ij) = +1$ , then  $a_{ij}^\sigma = 1$  and if  $e = ij$  is an edge and  $\sigma(ij) = -1$ , then  $a_{ij}^\sigma = -1$ , and if  $e = ij$  is not an edge, then  $a_{ij}^\sigma = 0$ . The adjacency matrix of a signed graph is symmetric (see [6] for some basic results on the adjacency spectrum of signed graphs). The signed Laplacian matrix of  $G^\sigma$  is the matrix  $L(G^\sigma) = D(G) - A(G^\sigma)$ , where  $D(G)$  is the diagonal matrix of vertex degrees. Let  $G$  be a graph with adjacency matrix  $A(G)$ . We say that  $\lambda(G)$  is an *eigenvalue of  $G$*  if there exists a non-zero vector  $X$  such that  $A(G)X = \lambda(G)X$ .

Throughout,  $t$  is the number of triangles,  $t^+$  (resp.,  $t^-$ ) is the number of balanced (resp., unbalanced) triangles,  $q$  is the number of quadrangles,  $q^+$  (resp.,  $q^-$ ) is the number of balanced (resp., unbalanced) quadrangles,  $t_i^+$  (resp.,  $t_i^-$ ) is the number of balanced (resp., unbalanced) triangles that are contain the  $i^{th}$  vertex,  $q_i^+$  (resp.,  $q_i^-$ ) is the number of balanced (resp., unbalanced) quadrangles that are contain the  $i^{th}$  vertex,  $t_{ij}^+$  (resp.,  $t_{ij}^-$ ) is the number of balanced (resp., unbalanced) triangles at edge  $ij$ ,  $U_n^+$  (resp.,  $U_n^-$ ) is a balanced graph obtained from  $C_{n-1}^+$  (resp.,  $C_{n-1}^-$ ) by attaching a leaf to one of its vertices, and  $B_5, B_4$  are two graphs obtained from two cycles  $C_3, C'_3$  of length 3 by identifying one vertex of  $C_3$  with one vertex of  $C'_3$  and identifying one edge of  $C_3$  with one edge of  $C'_3$ , respectively.  $|S_n|$  is the number of star graphs of order  $n$  and  $|P_n|$  is the number of paths of order  $n$ .

## 2. MAIN RESULTS

In this section, we would like to obtain  $k$ -th spectral moments and  $k$ -th Laplacian spectral moments of the signed generalized Petersen graphs. Taghvaei and Ashrafi in [3] and [4] determined the spectral moments and Laplacian spectral moments of generalized Petersen graph.

For the sake of completeness we mention below three important lemmas which are crucial in our main results in this paper (see, [1, 5]).

**Lemma 2.1.** *Let  $G^\sigma$  be a signed graph with signed adjacency matrix  $A(G^\sigma)$ . Then the  $(i, j)$ -entry of the matrix  $A^k(G^\sigma)$  is equal to  $w_{ij}^+(k) - w_{ij}^-(k)$ .*

Suppose that  $G^\sigma$  is a signed graph with adjacency matrix  $A(G^\sigma)$  and  $\lambda_1(G^\sigma), \lambda_2(G^\sigma), \dots, \lambda_n(G^\sigma)$  are eigenvalues of  $G^\sigma$  in non-increasing

order. Then,  $S_k(G^\sigma) = \sum_{i=1}^n \lambda_i^k(G^\sigma)$ ,  $k \geq 0$  is the  $k$ -th signed spectral moment of  $G^\sigma$ .

**Lemma 2.2.** *The  $k$ -th signed spectral moments of  $G^\sigma$  is equal to the number of closed walks of length  $k$ .*

**Lemma 2.3.** *Let  $G^\sigma$  be a signed graph with  $n$  vertices and  $m$  edges. Then,*

$$S_2(G^\sigma) = 2m \quad \text{and} \quad S_3(G^\sigma) = 6(t^+ - t^-).$$

It is also clear that  $S_0(G^\sigma) = n$  and  $S_1(G^\sigma) = 0$ .

**Theorem 2.4.** *Let  $G^\sigma$  be a signed graph with  $n$  vertices and  $m$  edges. Then, we have:*

$$\begin{aligned} S_4(G^\sigma) &= -2m + 2 \sum_{i=1}^n d_i^2 + 8(q^+ - q^-), \\ S_5(G^\sigma) &= 30(t^+ - t^-) + 10(U_4^+ - U_4^-) + 10(C_5^+ - C_5^-), \\ S_6(G^\sigma) &= -10m + 6 \sum_{i=1}^n d_i^2 + 6|P_4| + 12|S_4| + 12(U_5^+ - U_5^-) \\ &\quad + 36(B_4^+ - B_4^-) + 24(B_5^+ - B_5^-) + 24t(G) + 48(q^+ - q^-) \\ &\quad + 12(C_6^+ - C_6^-). \end{aligned}$$

*Proof.* Vertices that belong to a closed walk of length 4 induce subgraph in  $G^\sigma$  are isomorphic to  $P_2, P_3$  and  $q$ . Now, in the signed graph  $G^\sigma$ , the number of closed walks of length 4 which span these subgraphs is equal to  $2|P_2|, 4|P_3|, 8(q^+ - q^-)$ , respectively. Vertices that belong to a closed walk of length 5 induce subgraph in  $G^\sigma$  are isomorphic to  $t, U_4$  and  $C_5$ . In the signed graph  $G^\sigma$ , the number of closed walks of length 5 which span these subgraphs is equal to  $30(t^+ - t^-), 10(U_4^+ - U_4^-)$  and  $10(C_5^+ - C_5^-)$ , respectively. Now, we compute  $S_6(G^\sigma)$ . Vertices that belong to a closed walk of length 6 induce subgraph in  $G^\sigma$  are isomorphic to  $P_2, P_3, P_4, S_4, U_5, B_4, B_5, t, q, C_6$ . In the signed graph  $G^\sigma$  the number of closed walks of length 6 which span these subgraphs is equal to  $2|P_2|, 12|P_3|, 6|P_4|, 12|S_4|, 12(U_5^+ - U_5^-), 36(B_4^+ - B_4^-), 24(B_5^+ - B_5^-), 24t, 48(q^+ - q^-)$  and  $12(C_6^+ - C_6^-)$ , respectively. Note that the number of 3-path in connected graph  $G$  are equal to  $-m + \frac{1}{2} \sum_{i=1}^n d_i^2$ .  $\square$

Let  $T_k(G^\sigma) = \sum_{i=1}^n \mu_i^k(G^\sigma)$ ,  $k \geq 0$ , be the  $k$ -th spectral moment for the Laplacian spectrum of a signed graph  $G^\sigma$ , where  $\mu_1(G^\sigma), \mu_2(G^\sigma), \dots, \mu_n(G^\sigma)$  are the Laplacian eigenvalues. Belardo [1] proved that:

**Theorem 2.5.** [1] *Let  $G^\sigma$  be a signed graph with  $n$  vertices,  $m$  edges and degree sequence  $(d_1, \dots, d_n)$ . Then we have:*

$$\begin{aligned} T_0(G^\sigma) &= n, \\ T_1(G^\sigma) &= \sum_{i=1}^n d_i = 2m, \\ T_2(G^\sigma) &= 2m + \sum_{i=1}^n d_i^2, \\ T_3(G^\sigma) &= 6(t^- - t^+) + 3 \sum_{i=1}^n d_i^2 + \sum_{i=1}^n d_i^3. \end{aligned}$$

**Theorem 2.6.** *Let  $G^\sigma$  be a signed graph with  $n$  vertices,  $m$  edges and degree sequence  $(d_1, \dots, d_n)$ . Then we have:*

$$\begin{aligned} T_4(G^\sigma) &= S_4(G^\sigma) + 4 \sum_{i=1}^n d_i^3 + \sum_{i=1}^n d_i^4 + 4 \sum_{v_i \sim v_j}^m d_i d_j - 8 \sum_{i=1}^n d_i (t_i^+ - t_i^-), \\ T_5(G^\sigma) &= -S_5(G^\sigma) - 5 \sum_{i=1}^n d_i^2 + 5 \sum_{i=1}^n d_i^3 + 5 \sum_{i=1}^n d_i^4 + \sum_{i=1}^n d_i^5 \\ &+ 10 \left( \sum_{v_i \sim v_j}^m d_i d_j + \frac{1}{2} \sum_{i=1}^n \sum_{i \neq j}^n d_i^2 d_j a_{ij}^\sigma - \sum_{i=1}^n d_i^2 (t_i^+ - t_i^-) \right. \\ &\left. + \sum_{i=1}^n d_i (q_i^+ - q_i^-) - \sum_{v_i \sim v_j}^m d_i d_j (t_{ij}^+ - t_{ij}^-) \right), \end{aligned}$$

where  $d_i$  is the degree of  $i$ -th vertex.

*Proof.* Let  $L(G^\sigma) = D(G) - A(G^\sigma)$  be the Laplacian matrix of a signed graph  $G^\sigma$  and  $A(G^\sigma) = A^\sigma$ . First, we obtain  $T_4(G^\sigma)$ . We have

$$\begin{aligned} T_4(G^\sigma) &= \sum_{i=1}^n \mu_i^4(G^\sigma) = \text{tr}(D - A^\sigma)^4 \\ &= \text{tr}((D - A^\sigma)^2(D - A^\sigma)^2) \\ &= \text{tr}(D^4) - 4\text{tr}(D^3 A^\sigma) - 4\text{tr}(D(A^\sigma)^3) + 4\text{tr}((A^\sigma)^2 D^2) \\ &+ 2\text{tr}(D A^\sigma D A^\sigma) + \text{tr}((A^\sigma)^4). \end{aligned}$$

By Theorem 2.4, we have  $tr((A^\sigma)^4) = S_4(G^\sigma)$ , where  $S_4(G^\sigma)$  is 4-th spectral moments of  $G^\sigma$ . We have,  $tr(D^3A^\sigma) = 0$ ,  $tr(D^4) = \sum_{i=1}^n d_i^4$ ,  $tr((A^\sigma)^2D^2) = \sum_{i=1}^n d_i^3$ ,  $tr(D(A^\sigma)^3) = 2\sum_{i=1}^n d_i(t_i^+ - t_i^-)$ . By direct computation, we can see that  $tr(DA^\sigma DA^\sigma) = 2\sum_{v_i \sim v_j}^m d_i d_j$ , where  $v_i \sim v_j$  is an edge of  $G^\sigma$ . This shows that,

$$T_4(G^\sigma) = \sum_{i=1}^n d_i^4 - 8\sum_{i=1}^n d_i(t_i^+ - t_i^-) + 4\sum_{i=1}^n d_i^3 + 4\sum_{v_i \sim v_j}^m d_i d_j + S_4(G^\sigma).$$

In a similar way, we can compute  $T_5(G^\sigma)$ . We have,

$$\begin{aligned} T_5(G^\sigma) &= \sum_{i=1}^n \mu_i^5(G^\sigma) = tr(D - A^\sigma)^5 \\ &= tr(D^5) + 5tr(D^3(A^\sigma)^2) - 5tr(D^2(A^\sigma)^3) + 5tr(D^2A^\sigma DA^\sigma) \\ &\quad - 5tr(D(A^\sigma)^2DA^\sigma) + 5tr(D(A^\sigma)^4) - tr((A^\sigma)^5). \end{aligned}$$

By Theorem 2.4, we get  $tr((A^\sigma)^5) = S_5(G^\sigma)$ , where  $S_5(G^\sigma)$  is 5-th spectral moments of  $G^\sigma$ . We have,  $tr(D^5) = \sum_{i=1}^n d_i^5$ ,  $tr(D^3(A^\sigma)^2) = \sum_{i=1}^n d_i^4$ ,

$$tr(D^2(A^\sigma)^3) = 2\sum_{i=1}^n d_i^2(t_i^+ - t_i^-), \quad tr(D(A^\sigma)^4) = \sum_{i=1}^n d_i^3 + 2\sum_{v_i \sim v_j}^m d_i d_j -$$

$$\sum_{i=1}^n d_i^2 + 2\sum_{i=1}^n d_i(q_i^+ - q_i^-), \quad \text{where } v_i \sim v_j \text{ is an edge of } G^\sigma, \quad tr(D(A^\sigma)^2DA^\sigma) =$$

$$2\sum_{v_i \sim v_j}^m d_i d_j(t_{ij}^+ - t_{ij}^-) \quad \text{and } tr(D^2A^\sigma DA^\sigma) = \sum_{i=1}^n \sum_{i \neq j}^n d_i^2 d_j a_{ij}^\sigma, \quad \text{where } a_{ij}^\sigma \text{ is an}$$

element of the signed adjacency matrix. This implies that,

$$\begin{aligned} T_5(G^\sigma) &= \sum_{i=1}^n d_i^5 + 5\sum_{i=1}^n d_i^4 - 10\sum_{i=1}^n d_i^2(t_i^+ - t_i^-) + 5\sum_{i=1}^n \sum_{i \neq j}^n d_i^2 d_j a_{ij}^\sigma \\ &\quad - 10\sum_{v_i \sim v_j}^m d_i d_j(t_{ij}^+ - t_{ij}^-) + 5\sum_{i=1}^n d_i^3 + 10\sum_{v_i \sim v_j}^m d_i d_j - 5\sum_{i=1}^n d_i^2 \\ &\quad + 10\sum_{i=1}^n d_i(q_i^+ - q_i^-) - S_5(G^\sigma). \end{aligned}$$

This completes the proof.  $\square$

**Corollary 2.7.** *Suppose that  $T_k(G^\sigma)$  and  $T_k(G)$  are the  $k$ -th Laplacian spectral moment of a signed graph  $G^\sigma$  and a simple graph  $G$ , respectively. Then we have:*

$$\begin{aligned} T_3(G^\sigma) &\leq T_3(G), \\ T_4(G^\sigma) &\leq T_4(G), \\ T_5(G^\sigma) &\leq T_5(G), \end{aligned}$$

with equality if and only if  $G^\sigma$  is balanced signed graph.

**Definition 2.8.** The signed generalized Petersen graph, denoted by  $GP^\sigma(n, k)$ , is a graph with vertices and edges given by

$$V(GP^\sigma(n, k)) = \{a_i, b_i | 1 \leq i \leq n\},$$

and

$$E(GP^\sigma(n, k)) = \{a_i b_i, a_i a_{i+1}, b_i b_{i+k} | 1 \leq i \leq n\},$$

respectively. Here,  $i + k$  are integers modulo  $n$ , where  $n > 6$ . Since  $GP(n, k) \cong GP(n, n - k)$ , suppose that  $k \leq \lfloor \frac{n-1}{2} \rfloor$ . Let  $A(n, k)$  and  $B(n, k)$  be the induced subgraphs of  $GP(n, k)$  with the set of vertices  $\{a_1, \dots, a_n\}$  and  $\{b_1, \dots, b_n\}$ , respectively. The subgraphs  $A(n, k)$  and  $B(n, k)$  are the outer and inner subgraphs of  $GP(n, k)$ , respectively.

**Theorem 2.9.** *If  $S_i(GP^\sigma(n, k))$ ,  $2 \leq i \leq 6$ , is a signed spectral moment of  $GP^\sigma(n, k)$ , then the following holds:*

$$\begin{aligned} S_2(GP^\sigma(n, k)) &= 6n, \\ S_3(GP^\sigma(n, k)) &= \begin{cases} 6(t^+ - t^-) & \text{if } 3|n, k = \frac{n}{3} \\ 0 & \text{otherwise,} \end{cases} \\ S_4(GP^\sigma(n, k)) &= \begin{cases} 30n + 8(q^+ - q^-) & \text{if } (4|n, k = \frac{n}{4}) \\ & \text{or } (k = 1) \\ 30n & \text{otherwise,} \end{cases} \\ S_5(GP^\sigma(n, k)) &= \begin{cases} 10(C_5^+ - C_5^-) & \text{if } (5|n, k = \frac{n}{5}) \\ & \text{or } (k = \frac{2n}{5}, n \neq 10) \\ & \text{or } (k = 2) \\ 60(t^+ - t^-) & \text{if } 3|n, k = \frac{n}{3} \\ 0 & \text{otherwise,} \end{cases} \end{aligned}$$

$$S_6(GP^\sigma(n, k)) = \begin{cases} 174n + 12(C_6^+ - C_6^-) & \text{if } (6|n, k = \frac{n}{6}, n \neq 18) \text{ or} \\ & (2|n, k = \frac{n}{2} - 1, n \neq 8) \text{ or} \\ & (k = 3, n \neq \{8, 9, 10, 12, 18\}) \\ 174n + 96(q^+ - q^-) & \text{if } 4|n, k = \frac{n}{4}, n \neq 12 \\ 174n + 96(q^+ - q^-) & \text{if } k = 1 \\ +12(C_6^+ - C_6^-) \\ 176n & \text{if } 3|n, k = \frac{n}{3} \\ 174n & \text{otherwise.} \end{cases}$$

*Proof.* By Theorem 2.3, we get  $S_2(GP^\sigma(n, k)) = 6n$ . Suppose that  $3|n$  and  $k = \frac{n}{3}$ . Then,  $b_i b_{i+\frac{n}{3}} b_{i+\frac{2n}{3}} b_i$  is a triangle of  $GP^\sigma(n, k)$ . By Theorem 2.3, we have  $S_3(GP^\sigma(n, \frac{n}{3})) = 6(t^+ - t^-)$ . Otherwise, there are no triangle in  $GP^\sigma(n, k)$  and also  $S_3(GP^\sigma(n, k)) = 0$ . Assume that  $4|n$  and  $k = \frac{n}{4}$ , then  $b_i b_{i+\frac{n}{4}} b_{i+\frac{2n}{4}} b_{i+\frac{3n}{4}} b_i$  is a quadrangle in  $GP^\sigma(n, k)$  and if  $k = 1$ , then  $a_i b_i b_{i+1} a_{i+1} a_i$  is a quadrangle in  $GP^\sigma(n, k)$ . Then, by Theorem 2.4, we have

$$\begin{aligned} S_4(GP^\sigma(n, \frac{n}{4})) &= -6n + 36n + 8(q^+ - q^-) \\ &= 30n + 8(q^+ - q^-). \end{aligned}$$

Otherwise,  $q^+ - q^- = 0$ . So,  $S_4(GP^\sigma(n, k)) = 30n$ . Assume that  $5|n$ ,  $k = \frac{n}{5}$  or  $k = \frac{2n}{5}$  and  $n \neq 10$ . If  $k = \frac{n}{5}$ , then  $b_i b_{i+\frac{n}{5}} b_{i+\frac{2n}{5}} b_{i+\frac{3n}{5}} b_{i+\frac{4n}{5}} b_i$  is a pentagon in  $GP^\sigma(n, k)$ , if  $k = \frac{2n}{5}$ , then  $b_i b_{i+\frac{n}{5}} b_{i+\frac{2n}{5}} b_{i+\frac{4n}{5}} b_{i+\frac{6n}{5}} b_{i+\frac{8n}{5}} b_i$  is another pentagon of  $GP^\sigma(n, k)$  and if  $k = 2$  and  $n \neq 10$ , then  $a_i b_i b_{i+2} a_{i+2} a_{i+1} a_i$  is a pentagon in  $GP^\sigma(n, k)$ . On the other hand,  $U_4^+ - U_4^- = 0 = t^+ - t^-$ . Thus,

$$S_5(GP^\sigma(n, \frac{n}{5})) = S_5(GP^\sigma(n, \frac{2n}{5})) = S_5(GP^\sigma(n, 2)) = 10(C_5^+ - C_5^-).$$

In other cases,  $S_5(GP^\sigma(n, k)) = 0$ . If  $3|n$ ,  $k = \frac{n}{3}$ , then we have  $U_4^+ - U_4^- = 3(t^+ - t^-)$  and  $C_5^+ - C_5^- = 0$ . Hence,

$$S_5(GP^\sigma(n, \frac{n}{3})) = 60(t^+ - t^-).$$

Now, we obtain  $S_6(GP^\sigma(n, k))$ . By Theorem 2.4, we have

$$\begin{aligned} S_6(G^\sigma) &= -10m + 6 \sum_{i=1}^n d_i^2 + 6|P_4| + 12|S_4| + 12(U_5^+ - U_5^-) \\ &\quad + 36(B_4^+ - B_4^-) + 24(B_5^+ - B_5^-) + 24t(G) + 48(q^+ - q^-) \\ &\quad + 12(C_6^+ - C_6^-). \end{aligned}$$

Moreover,  $|S_4| = 2n$  and  $B_4^+ - B_4^- = B_5^+ - B_5^- = 0$ . Now, assume that  $6|n$ ,  $k = \frac{n}{6}$  and  $n \neq 18$ . Then,  $b_i b_{i+\frac{n}{6}} b_{i+\frac{n}{3}} b_{i+\frac{n}{2}} b_{i+\frac{4n}{6}} b_{i+\frac{5n}{6}}$  is a hexagon in  $GP^\sigma(n, k)$ . If  $2|n$  and  $k = \frac{n}{2} - 1$ , then  $a_i b_i b_{i+\frac{n}{2}-1} b_{i+n-2} a_{i+n-2} a_{i+n-1} a_i$  is a hexagon of  $GP^\sigma(n, k)$  and if  $k = 3$  and  $n \neq \{8, 9, 10, 12, 18\}$ , then  $a_i b_i b_{i+3} a_{i+3} a_{i+2} a_{i+1} a_i$  is a hexagon in  $GP^\sigma(n, k)$ . Therefore,  $|P_4| = 12n$  and  $q^+ - q^- = 0 = U_5^+ - U_5^-$ . Thus,

$$\begin{aligned} S_6(GP^\sigma(n, \frac{n}{6})) &= S_6(GP^\sigma(n, 3)) = S_6(GP^\sigma(n, \frac{n}{2} - 1)) \\ &= -30n + 108n + 72n + 24n + 12(C_6^+ - C_6^-) \\ &= 174n + 12(C_6^+ - C_6^-). \end{aligned}$$

If  $3|n$  and  $k = \frac{n}{3}$ , then  $|P_4| = 11n$  and  $C_6^+ - C_6^- = 0$ . Hence,

$$\begin{aligned} S_6(GP^\sigma(n, \frac{n}{3})) &= -30n + 108n + 66n + 24n + 8n \\ &= 176n. \end{aligned}$$

Suppose that  $4|n$ ,  $k = \frac{n}{4}$  and  $n \neq 12$ . So  $U_5^+ - U_5^- = 4(q^+ - q^-)$ . Therefore,

$$\begin{aligned} S_6(GP^\sigma(n, \frac{n}{4})) &= -30n + 108n + 72n + 24n + 96(q^+ - q^-) \\ &= 174n + 96(q^+ - q^-). \end{aligned}$$

Finally if  $k = 1$ , then  $U_5^+ - U_5^- = 4(q^+ - q^-)$ . Since  $a_i a_{i+1} a_{i+2} b_{i+2} b_{i+4} b_i a_i$  is a hexagon in  $GP^\sigma(n, k)$ , we get

$$\begin{aligned} S_6(GP^\sigma(n, 1)) &= -30n + 108n + 72n + 24n + 96(q^+ - q^-) + 12(C_6^+ - C_6^-) \\ &= 174n + 96(q^+ - q^-) + 12(C_6^+ - C_6^-). \end{aligned}$$

Otherwise,  $C_6^+ - C_6^- = 0$  and  $|P_4| = 12n$ . Thus,

$$S_6(GP^\sigma(n, k)) = -30n + 108n + 72n + 24n = 174n.$$

This completes the proof.  $\square$

**Theorem 2.10.** *If  $T_i(GP^\sigma(n, k))$ ,  $0 \leq i \leq 5$ , is a signed Laplacian spectral moments of  $GP^\sigma(n, k)$ , then*

$$T_0(GP^\sigma(n, k)) = 2n, \quad T_1(GP^\sigma(n, k)) = 6n, \quad T_2(GP^\sigma(n, k)) = 24n,$$

$$\begin{aligned}
T_3(GP^\sigma(n, k)) &= \begin{cases} 108n - 6(t^+ - t^-) & \text{if } 3|n, k = \frac{n}{3} \\ 108n & \text{otherwise,} \end{cases} \\
T_4(GP^\sigma(n, k)) &= \begin{cases} 516n - 72(t^+ - t^-) & \text{if } 3|n, k = \frac{n}{3} \\ 516n + 8(q^+ - q^-) & \text{if } (4|n, k = \frac{n}{4}) \\ & \text{or } (k = 1) \\ 516n & \text{otherwise,} \end{cases} \\
T_5(GP^\sigma(n, k)) &= \begin{cases} 2556n - 10(C_5^+ - C_5^-) & \text{if } (5|n, k = \frac{n}{5} \text{ or } k = \frac{2n}{5}) \\ & \text{or } (k = 2) \\ 2556 + 120(q^+ - q^-) & \text{if } 4|n, k = \frac{n}{4} \\ 2556n - 600(t^+ - t^-) & \text{if } 3|n, k = \frac{n}{3} \\ 2556n + 30 \sum_{i=1}^{2n} (q^+ - q^-) & \text{if } k = 1 \\ 2556n & \text{otherwise.} \end{cases}
\end{aligned}$$

*Proof.* Since  $|V(GP^\sigma(n, k))| = 2n$  and  $|E(GP^\sigma(n, k))| = 3n$ , we get  $T_0(GP^\sigma(n, k)) = 2n$  and  $T_1(GP^\sigma(n, k)) = 6n$ . By Theorem 2.5, we have  $T_2(GP^\sigma(n, k)) = 6n + 18n = 24n$ . Now, consider  $T_3(GP^\sigma(n, k))$ . If  $3|n$  and  $k = \frac{n}{3}$ , then  $b_i b_{i+\frac{n}{3}} b_{i+\frac{2n}{3}}$  is a triangle of  $GP^\sigma(n, k)$ . So, the number of triangles in  $GP^\sigma(n, k)$  is either equal to  $t^+ - t^-$  or equal to zero. In the first case, we have:

$$\begin{aligned}
T_3(GP^\sigma(n, k)) &= -6(t^+ - t^-) + 54n + 54n \\
&= -6(t^+ - t^-) + 108n,
\end{aligned}$$

and in the second case, we have  $T_3(GP^\sigma(n, k)) = 54n + 54n = 108n$ . Now, we compute  $T_4(GP^\sigma(n, k))$ . If  $3|n$  and  $k = \frac{n}{3}$ , then the number of triangles containing the outer subgraph of  $GP^\sigma(n, k)$  is equal to 0 and the number of triangles containing the inner subgraph of  $GP^\sigma(n, k)$  is equal to  $t^+ - t^-$ . On the other hand,  $S_4(GP^\sigma(n, k)) = 30n$ . Then, we have

$$\begin{aligned}
T_4(GP^\sigma(n, \frac{n}{3})) &= 30n - 72(t^+ - t^-) + 216n + 162n + 108n \\
&= 516n - 72(t^+ - t^-).
\end{aligned}$$

Suppose that  $4|n$  and  $k = \frac{n}{4}$ . Now,  $b_i b_{i+\frac{n}{4}} b_{i+\frac{2n}{4}} b_{i+\frac{3n}{4}}$  is a quadrangle in  $GP^\sigma(n, k)$ . Using Theorem 2.9, we have

$$S_4(GP^\sigma(n, k)) = 30n + 8(q^+ - q^-).$$

Since  $t_i = 0$ , we get

$$\begin{aligned} T_4(GP^\sigma(n, \frac{n}{4})) &= 30n + 8(q^+ - q^-) + 216n + 162n + 108n \\ &= 516n + 8(q^+ - q^-). \end{aligned}$$

If  $k = 1$ , then  $a_i b_i b_{i+1} a_{i+1} a_i$  is a quadrangle in  $GP^\sigma(n, k)$ . On the other hand, by Theorem 2.9, we get  $S_4(GP^\sigma(n, k)) = 30n + 8(q^+ - q^-)$ . Therefore,

$$\begin{aligned} T_4(GP^\sigma(n, 1)) &= 30n + 8(q^+ - q^-) + 216n + 162n + 108n \\ &= 516n + 8(q^+ - q^-). \end{aligned}$$

Otherwise,  $q^+ - q^- = 0$ . Now, by Theorem 2.9, we get

$$S_4(GP^\sigma(n, k)) = 30n.$$

So,

$$\begin{aligned} T_4(GP^\sigma(n, k)) &= 30n + 216n + 162n + 108n \\ &= 516n. \end{aligned}$$

Now, we compute  $T_5(GP^\sigma(n, k))$ . Let  $5|n$ ,  $k = \frac{n}{5}$  or  $k = \frac{2n}{5}$ . So,  $b_i b_{i+\frac{n}{5}} b_{i+\frac{2n}{5}} b_{i+\frac{3n}{5}} b_{i+\frac{4n}{5}} b_i$  is a pentagon in  $GP^\sigma(n, k)$ . If  $k = \frac{2n}{5}$ , then  $b_i b_{i+\frac{n}{5}} b_{i+\frac{2n}{5}} b_{i+\frac{4n}{5}} b_{i+\frac{6n}{5}} b_{i+\frac{8n}{5}} b_i$  is another pentagon of  $GP^\sigma(n, k)$ . Thus,

$$\begin{aligned} T_5(GP^\sigma(n, \frac{n}{5})) &= T_5(GP^\sigma(n, \frac{2n}{5})) = -90n + 270n + 810n + 486n \\ &\quad - 10(C_5^+ - C_5^-) + 270n + 810n \\ &= 2556n - 10(C_5^+ - C_5^-). \end{aligned}$$

If  $k = 2$ , then  $a_i b_i b_{i+2} a_{i+2} a_{i+1} a_i$  is a pentagon in  $GP^\sigma(n, k)$  and  $t^+ - t^- = 0 = q^+ - q^-$ . This implies that,

$$\begin{aligned} T_5(GP^\sigma(n, 2)) &= -90n + 270n + 810n + 486n - 10(C_5^+ - C_5^-) \\ &\quad + 270n + 810n \\ &= 2556n - 10(C_5^+ - C_5^-). \end{aligned}$$

If  $4|n$  and  $k = \frac{n}{4}$ , then  $C_5^+ - C_5^- = 0 = t^+ - t^-$ . Hence,

$$\begin{aligned} T_5(GP^\sigma(n, \frac{n}{4})) &= -90n + 270n + 810n + 486n + 270n + 810n \\ &\quad + 120(q^+ - q^-) \\ &= 2556n + 120(q^+ - q^-). \end{aligned}$$

If  $3|n$  and  $k = \frac{n}{3}$ , then  $C_5^+ - C_5^- = 0 = q^+ - q^-$ . So,

$$\begin{aligned} T_5(GP^\sigma(n, \frac{n}{3})) &= -90n + 270n + 810n + 486n - 600(t^+ - t^-) \\ &\quad + 270n + 810n \\ &= 2556n - 600(t^+ - t^-). \end{aligned}$$

Finally, if  $k = 1$ , then we get

$$\begin{aligned} T_5(GP^\sigma(n, 1)) &= -90n + 270n + 810n + 486n + 270n + 810n \\ &\quad + 30 \sum_{i=1}^{2n} (q^+ - q^-) \\ &= 2556n + 30 \sum_{i=1}^{2n} (q^+ - q^-). \end{aligned}$$

Otherwise, we have

$$\begin{aligned} T_5(GP^\sigma(n, k)) &= -90n + 270n + 810n + 486n + 270n + 810n \\ &= 2556n. \end{aligned}$$

This completes the proof.  $\square$

**Theorem 2.11.** (*Newton's identity [2]*): Let  $\lambda_1, \lambda_2, \dots, \lambda_n$  be the roots of the polynomial

$$\phi(A, x) = x^n + a_1x^{n-1} + \dots + a_{n-1}x + a_n,$$

with signed spectral moment  $S_k$ . Then

$$a_k = -\frac{1}{k}(S_k + S_{k-1}a_1 + S_{k-2}a_2 + \dots + S_1a_{k-1}).$$

Now, by Newton's identity we compute coefficients of characteristic polynomial of signed generalized Petersen graphs and coefficients of Laplacian characteristic polynomial of signed generalized Petersen graphs.

**Lemma 2.12.** *Let  $a_i(G^\sigma)$ ,  $1 \leq i \leq 5$ , be the coefficients of characteristic polynomial of a signed graph  $G^\sigma$ . Then, we have:*

$$\begin{aligned} a_1(G^\sigma) &= 0, \\ a_2(G^\sigma) &= -m, \\ a_3(G^\sigma) &= -2(t^+ - t^-), \\ a_4(G^\sigma) &= \frac{1}{2}(m^2 + m - 4(q^+ - q^-) - \sum_{i=1}^n d_i^2), \\ a_5(G^\sigma) &= (2m - 6)(t^+ - t^-) - 2(U_4^+ - U_4^-) - 2(C_5^+ - C_5^-). \end{aligned}$$

*Proof.* Using Lemma 2.3, Theorem 2.4 and Theorem 2.11, we get the desired result.  $\square$

**Lemma 2.13.** *Let  $b_i(G^\sigma)$ , for  $1 \leq i \leq 4$ , be the coefficients of Laplacian characteristic polynomial of a signed graph  $G^\sigma$ . Then, we have:*

$$\begin{aligned} b_1(G^\sigma) &= -2m, \\ b_2(G^\sigma) &= -\frac{1}{2}(2m - 4m^2 + \sum_{i=1}^n d_i^2), \\ b_3(G^\sigma) &= -\frac{1}{3}(4m^3 - 6m^2 + 3(1 - m) \sum_{i=1}^n d_i^2 + \sum_{i=1}^n d_i^3 + 6(t^- - t^+)), \\ b_4(G^\sigma) &= -\frac{1}{4} \left( -2m - 2m^2 + 8m^3 - \frac{8m^4}{3} + (2 - 10m + 4m^2) \sum_{i=1}^n d_i^2 \right. \\ &\quad + \left. \left( 4 - \frac{8m}{3} \right) \sum_{i=1}^n d_i^3 + \sum_{i=1}^n d_i^4 + 4 \sum_{v_i \sim v_j} d_i d_j - 8 \sum_{i=1}^n d_i (t_i^+ - t_i^-) \right. \\ &\quad \left. + 8(q^+ - q^-) + 16m(t^+ - t^-) - \frac{1}{2} \left( \sum_{i=1}^n d_i^2 \right)^2 \right). \end{aligned}$$

*Proof.* Using Theorem 2.5 and Theorem 2.11, we get the desired result.  $\square$

**Corollary 2.14.** *Let  $a_i(GP^\sigma(n, k))$  for  $1 \leq i \leq 5$ , be the coefficients of characteristic polynomial of a signed generalized Petersen graph. Then,*

we have:

$$\begin{aligned}
a_1(GP^\sigma(n, k)) &= 0, \\
a_2(GP^\sigma(n, k)) &= -3n, \\
a_3(GP^\sigma(n, k)) &= \begin{cases} -2(t^+ - t^-) & \text{if } 3|n, k = \frac{n}{3} \\ 0 & \text{otherwise,} \end{cases} \\
a_4(GP^\sigma(n, k)) &= \begin{cases} \frac{1}{2}(9n^2 - 15n - 4(q^+ - q^-)) & \text{if } (4|n, k = \frac{n}{4}) \\ & \text{or } (k = 1) \\ \frac{1}{2}(9n^2 - 15n) & \text{otherwise,} \end{cases} \\
a_5(GP^\sigma(n, k)) &= \begin{cases} -2(C_5^+ - C_5^-) & \text{if } (5|n, k = \frac{n}{5} \text{ or } k = \frac{2n}{5}) \\ & \text{or } (k = 2) \\ (6n - 12)(t^+ - t^-) & \text{if } 3|n, k = \frac{n}{3} \\ 0 & \text{otherwise.} \end{cases}
\end{aligned}$$

*Proof.* Since  $|E(GP^\sigma(n, k))| = 3n$ , then we have  $a_2(GP^\sigma(n, k)) = -3n$ . Assume that  $3|n$  and  $k = \frac{n}{3}$ . Then,  $b_i b_{i+\frac{n}{3}} b_{i+\frac{2n}{3}} b_i$  is a triangle of  $GP^\sigma(n, k)$ . Thus,

$$a_3(GP^\sigma(n, \frac{n}{3})) = -2(t^+ - t^-).$$

Otherwise, there is no triangle in  $GP^\sigma(n, k)$  and  $a_3(GP^\sigma(n, k)) = 0$ . Suppose that  $4|n$  and  $k = \frac{n}{4}$ . Then,  $b_i b_{i+\frac{n}{4}} b_{i+\frac{2n}{4}} b_{i+\frac{3n}{4}} b_i$  is a quadrangle in  $GP^\sigma(n, k)$ . So,

$$\begin{aligned}
a_4(GP^\sigma(n, \frac{n}{4})) &= \frac{1}{2}(9n^2 + 3n - 4(q^+ - q^-) - 18n) \\
&= \frac{1}{2}(9n^2 - 4(q^+ - q^-) - 15n).
\end{aligned}$$

Otherwise,  $q^+ - q^- = 0$ . Thus,

$$\begin{aligned}
a_4(GP^\sigma(n, k)) &= \frac{1}{2}(9n^2 + 3n - 18n) \\
&= \frac{1}{2}(9n^2 - 15n).
\end{aligned}$$

Now, assume that  $k = 1$ . Then,  $a_i b_i b_{i+1} a_{i+1} a_i$  is a quadrangle in  $GP^\sigma(n, k)$ . Therefore,

$$\begin{aligned} a_4(GP^\sigma(n, 1)) &= \frac{1}{2}(9n^2 + 3n - 4(q^+ - q^-) - 18n) \\ &= \frac{1}{2}(9n^2 - 4(q^+ - q^-) - 15n). \end{aligned}$$

Suppose that  $5|n$ ,  $k = \frac{n}{5}$  or  $k = \frac{2n}{5}$ . If  $k = \frac{n}{5}$ , then  $b_i b_{i+\frac{n}{5}} b_{i+\frac{2n}{5}} b_{i+\frac{3n}{5}} b_{i+\frac{4n}{5}} b_i$  is a pentagon in  $GP^\sigma(n, k)$  and if  $k = \frac{2n}{5}$ , then  $b_i b_{i+\frac{n}{5}} b_{i+\frac{2n}{5}} b_{i+\frac{4n}{5}} b_{i+\frac{6n}{5}} b_{i+\frac{8n}{5}} b_i$  is an another pentagon of  $GP^\sigma(n, k)$ . So,  $U_4^+ - U_4^- = 0 = t^+ - t^-$ . This yields that,

$$\begin{aligned} a_5(GP^\sigma(n, \frac{n}{5})) &= a_5(GP^\sigma(n, \frac{2n}{5})) \\ &= -2(C_5^+ - C_5^-). \end{aligned}$$

If  $k = 2$ , then  $a_i b_i b_{i+2} a_{i+2} a_{i+1} a_i$  is a pentagon in  $GP^\sigma(n, k)$  and  $U_4^+ - U_4^- = 0 = t^+ - t^-$ . This implies that  $a_5(GP^\sigma(n, 2)) = -2(C_5^+ - C_5^-)$ . In other cases, we have  $a_5(GP^\sigma(n, k)) = 0$ . If  $3|n$ ,  $k = \frac{n}{3}$ , then  $U_4^+ - U_4^- = 3(t^+ - t^-) = n$  and  $C_5^+ - C_5^- = 0$ . Hence,

$$\begin{aligned} a_5(GP^\sigma(n, \frac{n}{3})) &= (2(3n) - 2(3) - 6)(t^+ - t^-) \\ &= (6n - 12)(t^+ - t^-). \end{aligned}$$

Otherwise,  $a_5(GP^\sigma(n, k)) = 0$ . This completes the proof.  $\square$

**Corollary 2.15.** *Let  $b_i(GP^\sigma(n, k))$ , for  $1 \leq i \leq 4$ , be the coefficients of Laplacian characteristic polynomial of a signed generalized Petersen*

graph. Then, we have:

$$\begin{aligned}
 b_1(GP^\sigma(n, k)) &= -6n, \\
 b_2(GP^\sigma(n, k)) &= 18n^2 - 24n, \\
 b_3(GP^\sigma(n, k)) &= \begin{cases} -36n^3 + 72n^2 - \frac{108n}{3} & \text{if } 3|n, k = \frac{n}{3} \\ +2(t^+ - t^-) & \\ -36n^3 + 72n^2 - \frac{108n}{3} & \text{otherwise,} \end{cases} \\
 b_4(GP^\sigma(n, k)) &= \begin{cases} 54n^4 - 216n^3 + 288n^2 - 129n & \text{if } (4|n, k = \frac{n}{4}) \\ -2(q^+ - q^-) & \text{or } (k = 1) \\ 54n^4 - 216n^3 + 288n^2 - 129n & \text{if } 3|n, k = \frac{n}{3} \\ -(12n - 18)(t^+ - t^-) & \\ 54n^4 - 216n^3 + 288n^2 - 129n & \text{otherwise.} \end{cases}
 \end{aligned}$$

*Proof.* Using Lemma 2.13 and a similar method as used in the proof of Corollary 2.14, one can see that the assertion holds.  $\square$

**Acknowledgment.** The research of the second author was supported by the University of Kashan (Grant No. 682410/10). We thank F. Taghvaei for all of assistances that improved this paper.

#### REFERENCES

1. F. Belardo and P. Petecki, Spectral characterizations of signed lollipop graphs, *Linear Algebra Appl.* **480** (2015) 144–167.
2. M. Ghorbani and E. Bani-Asadi, Remarks on characteristic coefficients of fullerene graphs, *Applied Math. Comput.* **230** (2014) 428–435.
3. F. Taghvaei, On the Spectral Moment of Graphs, *MSC Thesis, University of Kashan*, January 2013.
4. F. Taghvaei and A. R. Ashrafi, Ordering some regular graphs with respect to spectral moments, 2017, preprint.
5. Y. P. Wu and H. Q. Lin, Lexicographical ordering by spectral moments of trees with a prescribed diameter, *Linear Algebra Appl.* **433** (2010) 1707–1713.
6. T. Zaslavsky, Signed graphs, *Discrete Appl. Math.* **4** (1982) 47–74.

#### Elham Ghasemian

Department of Pure Mathematics, Faculty of Mathematical Sciences, University of Kashan, Kashan 87317-53153, I. R. Iran.

Email: e.ghasemian@grad.kashanu.ac.ir

**G. H. Fath-tabar**

Department of Pure Mathematics, Faculty of Mathematical Sciences, University of  
Kashan, Kashan 87317-53153, I. R. Iran.

Email: [fathtabar@kashanu.ac.ir](mailto:fathtabar@kashanu.ac.ir)

SIGNED GENERALIZED PETERSEN GRAPH AND ITS CHARACTERISTIC POLYNOMIAL

E. GHASEMIAN AND G.H. FATH-TABAR

گراف پترسن تعمیم یافته علامت دار و ضریب چندجمله ای مشخصه ای آن

الهام قاسمیان و غلام حسین فتح تبار  
دانشکده علوم ریاضی دانشگاه کاشان، ایران، کاشان

فرض کنید  $G^\sigma$  گراف علامت دار است که در آن  $G = (V, E)$  گراف ساده و  $\sigma : E(G) \rightarrow \{\pm\}$  تابع علامت روی مجموعه یال های  $E(G)$  است. در این مقاله  $k$ -امین گشتاور طیفی و  $k$ -امین گشتاور طیفی لاپلاسی گراف علامت دار را به همراه ضرایب چندجمله ای مشخصه و ضرایب چندجمله ای مشخصه لاپلاسی گراف های علامت دار به دست آوریم.

کلمات کلیدی: گراف علامت دار، گراف پترسن تعمیم یافته علامت دار، ماتریس مجاورت.