

## ON MAXIMAL IDEALS OF $\mathcal{R}_\infty L$

A.A. ESTAJI\* AND A. MAHMOUDI DARGHADAM

ABSTRACT. Let  $L$  be a completely regular frame and  $\mathcal{R}L$  be the ring of real-valued continuous functions on  $L$ . We consider the set

$\mathcal{R}_\infty L = \{\varphi \in \mathcal{R}L : \uparrow \varphi(\frac{-1}{n}, \frac{1}{n}) \text{ is a compact frame for any } n \in \mathbb{N}\}$ .

Suppose that  $C_\infty(X)$  is the family of all functions  $f \in C(X)$  for which the set  $\{x \in X : |f(x)| \geq \frac{1}{n}\}$  is compact, for every  $n \in \mathbb{N}$ . Kohls has shown that  $C_\infty(X)$  is precisely the intersection of all the free maximal ideals of  $C^*(X)$ . The aim of this paper is to extend this result to the real continuous functions on a frame and hence we show that  $\mathcal{R}_\infty L$  is precisely the intersection of all the free maximal ideals of  $\mathcal{R}^*L$ . This result is used to characterize the maximal ideals in  $\mathcal{R}_\infty L$ .

### 1. INTRODUCTION

We denote by  $C(X)$  ( $C^*(X)$ ) the ring of all (bounded) real-valued continuous functions on a space  $X$  which is a nonempty completely regular Hausdorff space.  $C_\infty(X)$ , the subring of all functions  $C(X)$  which vanish at infinity, was introduced by Kohls in [16] (also, see [2, 1, 3, 18, 20] for more details). He shows that:

**Proposition 1.1.** [16, Lemma 3.2] *The ring  $C_\infty(X)$  is the intersection of the free maximal ideals of  $C^*(X)$ .*

Azarpanah and Soundararajan in [4], show that  $C_\infty(X)$  is an ideal in  $C^*(X)$  but not in  $C(X)$ , see also [16] and 7D in [14]. In fact,  $C_\infty(X)$

---

MSC(2010): Primary: 06D22; Secondary: 06F25, 54C30

Keywords: Frame, Compact, Maximal ideal, Ring of real valued continuous functions.

Received: 26 September 2017, Accepted: 12 January 2018.

\*Corresponding author.

is the subring of  $C(X)$  and topological spaces  $X$  for which  $C_\infty(X)$  is the ideal of  $C(X)$  are characterized in [4].

$\mathcal{R}_\infty L$ , the family of all functions  $f \in \mathcal{R}L$  for which  $\uparrow \varphi(\frac{-1}{n}, \frac{1}{n})$  is compact for each  $n \in \mathbb{N}$ , was introduced by Dube in [6].

In this paper, we are trying to show that  $\mathcal{R}_\infty L$  is a subring of  $\mathcal{R}L$  and an ideal of  $\mathcal{R}^*L$  (see Propositions 3.4 and 3.5) and it is not an ideal of  $\mathcal{R}L$  (see Example 3.6). Also, we prove that if for every  $a \in L$ ,  $\downarrow a$  is a locally compact frame implies  $\mathcal{R}^*(\downarrow a) = \mathcal{R}(\downarrow a)$ , then  $\mathcal{R}_\infty L$  is an ideal of  $\mathcal{R}L$  (see Proposition 3.9). In Section 4, we prove that for every completely regular frame  $L$ , it is a compact frame if and only if  $\mathcal{R}L = \mathcal{R}^*L = \mathcal{R}_\infty L$  (see Proposition 4.4). In Section 5, we show that the ring  $\mathcal{R}_\infty L$  is the intersection of all the free maximal ideals in  $\mathcal{R}^*L$  (see Proposition 5.7). In the last section, we study maximal ideals in the ring  $\mathcal{R}_\infty L$  and we show that if  $L$  is a completely regular frame, then every maximal ideal of  $\mathcal{R}_\infty L$  is strongly fixed ideal (see Proposition 6.6). In fact,  $M$  is a maximal ideal of  $\mathcal{R}_\infty L$  if and only if there exists  $p \in pt(L)$  such that

- (1)  $M = M_p^* \cap \mathcal{R}_\infty L$ , and
- (2)  $p \in \uparrow \varphi(\frac{-1}{n}, \frac{1}{n})$ , for some  $\varphi \in \mathcal{R}_\infty L$  and  $n \in \mathbb{N}$ .

## 2. PRELIMINARIES

Regarding the frame of reals  $\mathcal{L}(\mathbb{R})$  and the  $f$ -ring  $\mathcal{R}L$  of continuous real-valued functions on frame  $L$ , we use the notations of [5]. The bounded part, in the  $f$ -ring sense, of  $\mathcal{R}L$  is denoted by  $\mathcal{R}^*L$  and is characterized by:

$$\varphi \in \mathcal{R}^*L \Leftrightarrow \varphi(p, q) = 1 \text{ for some } p, q \in \mathbb{Q}.$$

An element  $a$  of a frame  $L$  is said to be *rather below* an element  $b$ , written  $a \prec b$ , provided that  $a^* \vee b = \top$ . Also,  $a$  is *completely below*  $b$ , written  $a \prec\prec b$ , if there are elements  $(c_q)$  indexed by the rational numbers  $\mathbb{Q} \cap [0, 1]$  such that  $c_0 = a$ ,  $c_1 = b$ , and  $c_p \prec c_q$  for  $p < q$ . A frame  $L$  is said to be *regular* if  $a = \bigvee \{x \in L : x \prec a\}$  for each  $a \in L$ , and *completely regular* if  $a = \bigvee \{x \in L : x \prec\prec a\}$  for each  $a \in L$ .

An element  $p$  of  $L$  is *point* (or *prime*) whenever  $p < \top$  and  $a \wedge b \leq p$  implies that  $a \leq p$  or  $b \leq p$ . We denote the set of all points of  $L$  by  $pt(L)$  or  $\Sigma L$ .

An ideal  $J$  of  $L$  is completely regular, if for each  $x \in J$  there exists  $y \in J$  such that  $x \prec\prec y$ . The Stone-Ćech compactification of  $L$  is the frame  $\beta L$  consisting of completely regular ideals of  $L$  together with the dense onto frame homomorphism  $j_L : \beta L \rightarrow L$  given by join. We denote

the right adjoint of  $j_L$  by  $r_L$ , and recall that  $r_L(a) = \{x \in L : x \ll a\}$ , for all  $a \in L$ .

Let  $L$  be a frame,  $a \in L$  and  $\alpha \in \mathcal{R}L$ . The sets  $\{r \in \mathbb{Q} : \alpha(-, r) \leq a\}$  and  $\{s \in \mathbb{Q} : \alpha(s, -) \leq a\}$ , are denoted by  $L(a, \alpha)$  and  $U(a, \alpha)$  respectively. For  $a \neq \top$ , it is obvious that  $r \leq s$ , for each  $r \in L(a, \alpha)$  and  $s \in U(a, \alpha)$ . In fact, we have:

**Proposition 2.1.** [8] *Let  $L$  be a frame and  $p$  be a prime element of  $L$ . There exists a unique map  $\tilde{p} : \mathcal{R}L \rightarrow \mathbb{R}$  such that  $r \leq \tilde{p}(\alpha) \leq s$ , for each  $\alpha \in \mathcal{R}L$ ,  $r \in L(p, \alpha)$  and  $s \in U(p, \alpha)$ .*

**Proposition 2.2.** [8] *If  $p$  is a prime element of a frame  $L$ , then  $\tilde{p} : \mathcal{R}L \rightarrow \mathbb{R}$  is an onto  $f$ -ring homomorphism.*

Let  $\alpha \in \mathcal{R}L$ . We define  $\alpha[p] = \tilde{p}(\alpha)$  for all  $p \in \Sigma L$ , and define

$$Z(\alpha) = \{p \in \Sigma L : \alpha[p] = 0\}.$$

This set is said to be a zero-set in  $L$  (see [11]). For  $A \subseteq \mathcal{R}L$ , we write  $Z[A]$  to designate the family of zero-sets  $\{Z(\alpha) : \alpha \in A\}$ . The family  $Z[\mathcal{R}L]$  of all zero-sets in  $L$  will also be denoted, for simplicity, by  $Z[L]$  (also, see [10, 12, 15] for more details on the zero-sets and their application in  $\mathcal{R}L$ ). For undefined terms and notations, the readers are referred to [9, 17].

### 3. TOPICS IN $\mathcal{R}_\infty L$ IS AN IDEAL OF $\mathcal{R}L$ AND AN IDEAL OF $\mathcal{R}^*L$

The following lemma is proved in [6]. It will be used in this paper.

**Lemma 3.1.** *For every  $a, b \in L$ , if  $\uparrow a$  and  $\uparrow b$  are compact, then  $\uparrow(a \wedge b)$  is compact.*

*Remark 3.2.* For every  $a, b \in L$ , if  $\uparrow a$  is compact and  $a \leq b$ , then  $\uparrow b$  is compact.

*Remark 3.3.* Consider  $\varphi \in \mathcal{R}_\infty L$  and  $0 < \varepsilon \in \mathbb{Q}$ . Then, there exists  $n \in \mathbb{N}$  such that  $\frac{1}{n} \leq \varepsilon$ . Since  $\varphi(\frac{-1}{n}, \frac{1}{n}) \leq \varphi(-\varepsilon, \varepsilon)$ , we can conclude from the Remark 3.2 that  $\uparrow \varphi(-\varepsilon, \varepsilon)$  is compact. Therefore, for every  $\varphi \in \mathcal{R}L$ ,  $\varphi \in \mathcal{R}_\infty L$  if and only if for every  $0 < \varepsilon \in \mathbb{Q}$ ,  $\uparrow \varphi(-\varepsilon, \varepsilon)$  is compact.

For every  $p, q, u, v \in \mathbb{Q}$ , we put

$$\langle p, q \rangle := \{r \in \mathbb{Q} : p < r < q\}$$

and

$$\langle p, q \rangle \langle u, v \rangle := \{rs : p < r < q, u < s < v\}.$$

In this paper, a subring of a commutative ring with identity does not imply the identity must belong to the subring.

**Proposition 3.4.**  $\mathcal{R}_\infty L$  is a subring of  $\mathcal{R}L$ .

*Proof.* Consider  $\varphi, \psi \in \mathcal{R}_\infty L$  and  $n \in \mathbb{N}$ . Since  $\uparrow \varphi(\frac{-1}{2n}, \frac{1}{2n})$  and  $\uparrow \psi(\frac{-1}{2n}, \frac{1}{2n})$  are compact frames, we can conclude from the Lemma 3.1 that  $\uparrow (\varphi(\frac{-1}{2n}, \frac{1}{2n}) \wedge \psi(\frac{-1}{2n}, \frac{1}{2n}))$  is a compact frame. The fact that

$$\varphi(\frac{-1}{2n}, \frac{1}{2n}) \wedge \psi(\frac{-1}{2n}, \frac{1}{2n}) \leq (\varphi + \psi)(\frac{-1}{n}, \frac{1}{n})$$

enables us to conclude at once that  $\uparrow (\varphi + \psi)(\frac{-1}{n}, \frac{1}{n})$  is a compact frame, by Remark 3.2. Therefore,  $\varphi + \psi \in \mathcal{R}_\infty L$ .

Let  $m \in \mathbb{N}$  such that  $\frac{1}{m} \leq \frac{1}{\sqrt{n}}$ . Since  $\uparrow \varphi(\frac{-1}{m}, \frac{1}{m})$  and  $\uparrow \psi(\frac{-1}{m}, \frac{1}{m})$  are compact and

$$\varphi(\frac{-1}{m}, \frac{1}{m}) \wedge \psi(\frac{-1}{m}, \frac{1}{m}) \leq (\varphi\psi)(\frac{-1}{n}, \frac{1}{n}),$$

we can conclude from the Lemma 3.1 and the Remark 3.2 that  $\uparrow (\varphi\psi)(\frac{-1}{n}, \frac{1}{n})$  is compact. Hence,  $\varphi\psi \in \mathcal{R}_\infty L$ .  $\square$

**Proposition 3.5.**  $\mathcal{R}_\infty L$  is an ideal of  $\mathcal{R}^*L$ .

*Proof.* Consider  $\varphi \in \mathcal{R}_\infty L$  and  $n \in \mathbb{N}$ . Since for all  $m \in \mathbb{N}$ ,

$$\varphi(-m, m) \in \uparrow \varphi(\frac{-1}{n}, \frac{1}{n})$$

and

$$\top = \bigvee_{m \in \mathbb{N}} \varphi(-m, m),$$

we conclude that there are  $m_1, m_1, \dots, m_k \in \mathbb{N}$  such that

$$\top = \bigvee_{1 \leq i \leq k} \varphi(-m_i, m_i).$$

If  $m = \text{Max}\{m_1, m_2, \dots, m_k\}$  then  $\varphi(-m, m) = \top$ , that is  $\varphi \in \mathcal{R}^*L$ . Therefore,  $\mathcal{R}_\infty L \subseteq \mathcal{R}^*L$ .

Now, suppose that  $\varphi \in \mathcal{R}_\infty L$  and  $\psi \in \mathcal{R}^*L$ . It suffices to show that  $\varphi\psi \in \mathcal{R}_\infty L$ . There exists  $m \in \mathbb{N}$  such that  $\psi(-m, m) = \top$ , by hypothesis. Consider  $n \in \mathbb{N}$ . Since

$$\langle -\frac{1}{mn}, \frac{1}{mn} \rangle \langle -m, m \rangle \subseteq \langle -\frac{1}{n}, \frac{1}{n} \rangle,$$

we have

$$\varphi(-\frac{1}{mn}, \frac{1}{mn}) = \varphi(-\frac{1}{mn}, \frac{1}{mn}) \wedge \psi(-m, m) \leq (\varphi\psi)(-\frac{1}{n}, \frac{1}{n}).$$

Since  $\uparrow \varphi(-\frac{1}{mn}, \frac{1}{mn})$  is a compact frame, we can conclude from the Remark 3.2 that  $\uparrow (\varphi\psi)(-\frac{1}{n}, \frac{1}{n})$  is a compact frame, hence  $\varphi\psi \in \mathcal{R}_\infty L$ .  $\square$

The following example shows that  $\mathcal{R}_\infty L$  is not an ideal of  $\mathcal{R}L$  in general.

**Example 3.6.** We consider the function  $\alpha : \mathcal{L}\mathbb{R} \rightarrow \mathcal{P}(\mathbb{N})$  defined by

$$\alpha(p, q) = \{ n \in \mathbb{N} : p < \frac{1}{n} < q \},$$

for every  $p, q \in \mathbb{Q}$ . We claim that  $\alpha$  is a frame map. To prove this, we check the relations (R1)-(R4) to identities in  $\mathcal{P}(\mathbb{N})$  (see [5]).

(R1). For every  $p, q, r, s \in \mathbb{Q}$ , we have

$$\begin{aligned} \alpha(p, q) \wedge \alpha(r, s) &= \{ n \in \mathbb{N} : p < \frac{1}{n} < q \} \cap \{ n \in \mathbb{N} : r < \frac{1}{n} < s \} \\ &= \{ n \in \mathbb{N} : p \vee r < \frac{1}{n} < q \wedge s \} \\ &= \alpha(p \vee r, q \wedge s) \\ &= \alpha((p, q) \wedge (r, s)). \end{aligned}$$

(R2). For every  $p, q, r, s \in \mathbb{Q}$  with  $p \leq r < q \leq s$ , we have

$$\begin{aligned} \alpha(p, q) \vee \alpha(r, s) &= \{ n \in \mathbb{N} : p < \frac{1}{n} < q \} \cup \{ n \in \mathbb{N} : r < \frac{1}{n} < s \} \\ &= \{ n \in \mathbb{N} : p \wedge r < \frac{1}{n} < q \vee s \} \\ &= \{ n \in \mathbb{N} : p < \frac{1}{n} < s \} \\ &= \alpha(p, s). \end{aligned}$$

(R3). For every  $p, q \in \mathbb{Q}$ , we have

$$\begin{aligned} \bigvee_{p < r < s < q} \alpha(r, s) &= \bigcup_{p < r < s < q} \{ n \in \mathbb{N} : r < \frac{1}{n} < s \} \\ &= \{ n \in \mathbb{N} : p < \frac{1}{n} < q \} \\ &= \alpha(p, q). \end{aligned}$$

(R4). It is clear that

$$\mathbb{N} = \top_{\mathcal{P}(\mathbb{N})} = \alpha(0, 2) \leq \bigcup_{p, q \in \mathbb{Q}} \alpha(p, q) \leq \mathbb{N},$$

then  $\bigvee_{p, q \in \mathbb{Q}} \alpha(p, q) = \top_{\mathcal{P}(\mathbb{N})}$ . Therefore,  $\alpha \in \mathcal{R}(\mathcal{P}(\mathbb{N}))$ .

Since, for any  $n \in \mathbb{N}$ ,

$$\alpha\left(\frac{-1}{n}, \frac{1}{n}\right) = \{m \in \mathbb{N} : n < m\} = \{n+1, n+2, n+3, \dots\},$$

we infer that  $\uparrow \alpha\left(\frac{-1}{n}, \frac{1}{n}\right)$  is a finite frame and hence it is a compact frame. Hence,  $\alpha \in \mathcal{R}_\infty(\mathcal{P}(\mathbb{N}))$ . Since

$$\uparrow \mathbf{1}\left(\frac{-1}{n}, \frac{1}{n}\right) = \uparrow \perp = \mathcal{P}(\mathbb{N})$$

is not a compact frame, we conclude that  $\mathbf{1} \notin \mathcal{R}_\infty(\mathcal{P}(\mathbb{N}))$ . Since

$$\text{coz}(\alpha) = \alpha(-, 0) \vee \alpha(0, -) = \mathbb{N} = \top_{\mathcal{P}(\mathbb{N})},$$

we conclude that

- (1)  $\alpha$  is unit and  $\alpha \in \mathcal{R}_\infty(\mathcal{P}(\mathbb{N}))$ .
- (2)  $\mathcal{R}_\infty(\mathcal{P}(\mathbb{N})) \subsetneq \mathcal{R}(\mathcal{P}(\mathbb{N}))$ .
- (3)  $\mathcal{R}_\infty(\mathcal{P}(\mathbb{N}))$  is not an ideal of  $\mathcal{R}(\mathcal{P}(\mathbb{N}))$ .

Let  $L$  be a frame. We say that  $a$  is *way below*  $b$  (or *relatively compact with respect to*  $b$ ) and write  $a \ll b$  if for any  $S \subseteq L$  with  $b \leq \bigvee S$ , there exists a finite set  $F \subseteq S$  such that  $a \leq \bigvee F$ .

A frame  $L$  is called *continuous* (or *locally compact*) whenever, for each  $a \in L$ ,  $a = \bigvee_{x \ll a} x$ .

**Lemma 3.7.** *For every completely regular frame  $L$  and  $\varphi \in \mathcal{R}_\infty L$ ,  $\downarrow \text{coz}(\varphi)$  is a locally compact frame.*

*Proof.* Consider  $a \in \downarrow \text{coz}(\varphi)$ . Let  $x \prec a \wedge \varphi\left(\left(-, -\frac{1}{n}\right) \vee \left(\frac{1}{n}, -\right)\right)$

and  $S \subseteq L$  with  $a \wedge \varphi\left(\left(-, -\frac{1}{n}\right) \vee \left(\frac{1}{n}, -\right)\right) \leq \bigvee S$ . Then

$$\begin{aligned} \varphi\left(-\frac{1}{n}, \frac{1}{n}\right) &\leq \left(\varphi\left(\left(-, -\frac{1}{n}\right) \vee \left(\frac{1}{n}, -\right)\right)\right)^* \\ &\leq a^* \vee \left(\varphi\left(\left(-, -\frac{1}{n}\right) \vee \left(\frac{1}{n}, -\right)\right)\right)^* \\ &= \left(a \wedge \varphi\left(\left(-, -\frac{1}{n}\right) \vee \left(\frac{1}{n}, -\right)\right)\right)^* \\ &\leq x^*. \end{aligned}$$

Using  $\varphi \in \mathcal{R}_\infty L$ , we conclude from Remark 3.2 that  $\uparrow x^*$  is a compact frame.

Since

$$\top = x^* \vee \left(a \wedge \varphi\left(\left(-, -\frac{1}{n}\right) \vee \left(\frac{1}{n}, -\right)\right)\right) \leq x^* \vee \bigvee S,$$

we infer that there are  $s_1, \dots, s_k \in S$  such that  $\top = \bigvee_{i=1}^k (x^* \vee s_i)$ , which implies that  $x \leq \bigvee_{i=1}^k s_i$ . Hence, if  $x \prec a \wedge \varphi\left(\left(-, -\frac{1}{n}\right) \vee \left(\frac{1}{n}, -\right)\right)$ ,

then  $x \ll a \wedge \varphi((- , -\frac{1}{n}) \vee (\frac{1}{n}, -))$ , for every  $x \in L$ . Therefore, the complete regularity of  $L$  insures that

$$\begin{aligned} a &= a \wedge \text{coz}(\varphi) \\ &= \bigvee_{n \in \mathbb{N}} (a \wedge \varphi((- , -\frac{1}{n}) \vee (\frac{1}{n}, -))) \\ &= \bigvee_{n \in \mathbb{N}} \bigvee \{ x \in L : x \prec a \wedge \varphi((- , -\frac{1}{n}) \vee (\frac{1}{n}, -)) \} \\ &\leq \bigvee_{n \in \mathbb{N}} \bigvee \{ x \in L : x \ll a \wedge \varphi((- , -\frac{1}{n}) \vee (\frac{1}{n}, -)) \} \\ &\leq \bigvee_{\substack{x \in L, \\ x \ll a}} x \\ &\leq a, \end{aligned}$$

and this completes the proof.  $\square$

**Lemma 3.8.** *Let  $\alpha \in \mathcal{R}L$  and  $\rho_3 : \mathcal{L}(\mathbb{R}) \rightarrow \mathcal{L}(\mathbb{R})$  by  $\rho_3(p, q) = (p^3, q^3)$ . Then the following statements hold:*

- (1)  $\rho_3 \in \mathcal{R}(\mathcal{L}(\mathbb{R}))$ .
- (2)  $\rho_3^3 = \text{id}_{\mathcal{L}(\mathbb{R})}$ .
- (3)  $(\alpha \circ \rho_3)^3 = \alpha$ .
- (4)  $\text{coz}(\alpha \circ \rho_3) = \text{coz}(\alpha)$ .
- (5) If  $\alpha \in \mathcal{R}_\infty L$ , then  $\alpha \circ \rho_3 \in \mathcal{R}_\infty L$ .

*Proof.* By [13], to complete the proof it suffices to show that if  $\alpha \in \mathcal{R}_\infty L$ , then  $\alpha \circ \rho_3 \in \mathcal{R}_\infty L$ . Consider  $\alpha \in \mathcal{R}_\infty L$ . Since for every  $n \in \mathbb{N}$ ,  $\uparrow \alpha \circ \rho_3(-\frac{1}{n}, \frac{1}{n}) = \uparrow \alpha(-\frac{1}{n^3}, \frac{1}{n^3})$  is a compact frame, we conclude that  $\alpha \circ \rho_3 \in \mathcal{R}_\infty L$ .  $\square$

**Proposition 3.9.** *Let  $L$  be a completely regular frame and for every  $a \in L$ , if  $\downarrow a$  is a locally compact frame, then  $\mathcal{R}^*(\downarrow a) = \mathcal{R}(\downarrow a)$ . Then  $\mathcal{R}_\infty L$  is an ideal of  $\mathcal{R}L$ .*

*Proof.* Consider  $\alpha \in \mathcal{R}L$  and  $\beta \in \mathcal{R}_\infty L$ . We put  $\beta^{\frac{1}{3}} = \beta \circ \rho_3$ . By Lemma 3.8, we have  $\alpha\beta^{\frac{1}{3}} \in \mathcal{R}L$ , which implies that  $\bar{\alpha} : \mathcal{L}\mathbb{R} \rightarrow \downarrow \text{coz}(\beta)$  given by  $\bar{\alpha}(u) = \alpha\beta^{\frac{1}{3}}(u) \wedge \text{coz}(\beta)$  is an element of  $\mathcal{R}(\downarrow \text{coz}(\beta))$ . Since, by Lemma 3.7,  $\downarrow \text{coz}(\beta)$  is a locally compact frame, we conclude that there exists  $n \in \mathbb{N}$  such that

$$\alpha\beta^{\frac{1}{3}}((- , -n) \vee (n, -)) \wedge \text{coz}(\beta) = \bar{\alpha}((- , -n) \vee (n, -)) = \perp.$$

By

$$\alpha\beta^{\frac{1}{3}}((- , -n) \vee (n, -)) \leq \text{coz}(\alpha\beta^{\frac{1}{3}}) \leq \text{coz}(\beta),$$

we infer that

$$\alpha\beta^{\frac{1}{3}}((- , -n) \vee (n, -)) = \perp,$$

which follows that  $\alpha\beta^{\frac{1}{3}} \in \mathcal{R}^*L$ . Since, by Lemma 3.8,  $\beta^{\frac{1}{3}} \in \mathcal{R}_\infty L$ , we conclude from Proposition 3.5 and Lemma 3.8 that  $\alpha\beta = \alpha\beta^{\frac{1}{3}}(\beta^{\frac{1}{3}})^2 \in \mathcal{R}_\infty L$  and this completes the proof.  $\square$

4. WHEN IS  $\mathcal{R}_\infty L$  EQUAL TO  $\mathcal{R}L$ ?

In this section, we characterize frames  $L$  for which  $\mathcal{R}_\infty L = \mathcal{R}L$ . Let  $I$  be an ideal in  $\mathcal{R}L$  or  $\mathcal{R}^*L$ . If  $\bigvee\{\text{coz}(\varphi) : \varphi \in I\} < \top$ , we call  $I$  a fixed ideal; if  $\bigvee\{\text{coz}(\varphi) : \varphi \in I\} = \top$ , then  $I$  is a free ideal.

**Lemma 4.1.** *If  $I$  is a free ideal in  $\mathcal{R}L$  and  $a \in \text{Coz}(L)$  is a compact element of  $\text{Coz}(L)$ , then there exists  $\varphi \in I$  such that  $a = \text{coz}(\varphi)$ .*

*Proof.* Evidently

$$a = a \wedge \top = \bigvee\{a \wedge \text{coz}(\varphi) : \varphi \in I\},$$

it follows that there are  $\varphi_1, \dots, \varphi_n \in I$  such that

$$a = a \wedge \bigvee_{i=1}^n \text{coz}(\varphi_i) = a \wedge \text{coz}(\varphi_1^2 + \dots + \varphi_n^2).$$

Since  $\text{Coz}(I)$  is an ideal of  $\text{Coz}(L)$  and

$$a \leq \text{coz}(\varphi_1^2 + \dots + \varphi_n^2) \in \text{Coz}(I)$$

we include that there exists  $\varphi \in I$  such that  $a = \text{coz}(\varphi)$ .  $\square$

**Corollary 4.2.** *The set*

$$\{a \in \text{Coz}(L) : a \text{ is a compact element of } \text{Coz}(L)\}$$

*is a subset of*

$$\bigcap\{\text{Coz}(I) : I \text{ is a free ideal in } \mathcal{R}L\}.$$

*Proof.* By Lemma 4.1, it is clear.  $\square$

The following proposition is proved by Dube in [6, Lemma 4.7], but here, in the proof of this proposition, a different approach is used.

**Proposition 4.3.** *For every completely regular frame  $L$ , the following statements are equivalent:*

- (1)  $L$  is a compact frame;
- (2) Every proper ideal  $I$  in  $\mathcal{R}L$  is fixed;
- (3) Every maximal ideal  $I$  in  $\mathcal{R}L$  is fixed.

*Proof.* (1)  $\Rightarrow$  (2). Let  $I$  be a proper free ideal in  $\mathcal{R}L$ , then by Lemma 4.1, there exists  $\varphi \in I$  such that  $\top = \text{coz}(\varphi)$ . It then follows that  $I$  contains a unit element. Hence,  $I = \mathcal{R}L$  and this is a contradiction.

(2)  $\Rightarrow$  (3). It is clear.

(3)  $\Rightarrow$  (1). Let  $\{a_\lambda\}_{\lambda \in \Lambda} \subseteq L$  such that  $\top = \bigvee_{\lambda \in \Lambda} a_\lambda$ . It is clear that

$$I = \{\varphi \in \mathcal{R}L : \exists \Lambda' \subseteq \Lambda (|\Lambda'| < \infty, \text{coz}(\varphi) \leq \bigvee_{\lambda \in \Lambda'} a_\lambda)\}$$

is an ideal of  $\mathcal{R}L$ . If  $I \neq \mathcal{R}L$ , then there exists a maximal ideal  $M$  such that  $I \subseteq M$ . Since  $L$  is completely regular frame, we infer that

$$\top = \bigvee_{\lambda \in \Lambda} a_\lambda = \bigvee \text{Coz}(I) \leq \bigvee \text{Coz}(M),$$

i.e.,  $\top = \bigvee \text{Coz}(M)$ , which is a contradiction. Now, we can assume that  $I = \mathcal{R}L$ . Then there exists  $\Lambda' \subseteq \Lambda$  such that  $|\Lambda'| < \infty$  and

$$\top = \text{coz}(\mathbf{1}) = \bigvee_{\lambda \in \Lambda'} a_\lambda,$$

this completes the proof of the proposition.  $\square$

**Proposition 4.4.** *For every completely regular frame  $L$ , then  $L$  is a compact frame if and only if  $\mathcal{R}L = \mathcal{R}^*L = \mathcal{R}_\infty L$ .*

*Proof. Necessity.*

Consider  $\varphi \in \mathcal{R}L$ ,  $n \in \mathbb{N}$  and  $a = \varphi(-\frac{1}{n}, \frac{1}{n})$ . Since  $L = \uparrow \perp$  is a compact frame and  $\perp \leq a$ , we can conclude from the Remark 3.2 that  $\uparrow a$  is a compact frame, i.e.,  $\varphi \in \mathcal{R}_\infty L$ .

*Sufficiency.* Since  $\mathbf{1} \in \mathcal{R}_\infty L$ , we infer that

$$L = \uparrow \perp = \uparrow \mathbf{1}(-1, 1)$$

is a compact frame.  $\square$

## 5. INTERSECTION OF FREE MAXIMAL IDEALS

In [16, Lemma 3.2], the intersection of the free maximal ideals in  $C^*(X)$  was characterized as the set of all functions that vanish at infinity (that is all functions  $f \in C(X)$  such that  $\{x : |f(x)| \geq \frac{1}{n}\}$  is compact for all  $n \in \mathbb{N}$ ). In this section, we show that this is also true for  $\mathcal{R}^*(L)$ .

**Proposition 5.1.** *If  $I$  is a proper free ideal in  $\mathcal{R}L$ , then*

$$\varphi(-\frac{1}{n}, \frac{1}{n}) \notin \text{Coz}(I),$$

for every  $\varphi \in \mathcal{R}_\infty L$  and  $n \in \mathbb{N}$ .

*Proof.* Consider  $\varphi \in \mathcal{R}L$  and  $n \in \mathbb{N}$ . Then

$$\top = \bigvee I = \bigvee \{\text{coz}(\alpha) \vee \varphi(-\frac{1}{n}, \frac{1}{n}) : \alpha \in I\}$$

and since  $\uparrow \varphi(-\frac{1}{n}, \frac{1}{n})$  is compact, we conclude that there are  $\alpha_1, \dots, \alpha_k \in I$  such that

$$\top = \left( \bigvee_{i=1}^k \text{coz}(\alpha_i) \right) \vee \varphi(-\frac{1}{n}, \frac{1}{n}) = \text{coz} \left( \sum_{i=1}^k \alpha_i^2 \right) \vee \varphi(-\frac{1}{n}, \frac{1}{n})$$

and  $\sum_{i=1}^k \alpha_i^2 \in I$ . If  $\varphi(-\frac{1}{n}, \frac{1}{n}) \in \text{Coz}(I)$ , then  $\top \in \text{Coz}(I)$ , i.e.,  $I = \mathcal{R}L$ , which is a contradiction. Hence,  $\varphi(-\frac{1}{n}, \frac{1}{n}) \notin \text{Coz}(I)$ .  $\square$

It is well known that  $\mathfrak{t}_L : \mathcal{R}(\beta L) \rightarrow \mathcal{R}^*L$  given by  $\mathfrak{t}_L(\alpha) = j_L\alpha$  is the ring isomorphism. Also, we will denote  $\varphi^\beta = \mathfrak{t}_L^{-1}(\varphi)$ , for every  $\varphi \in \mathcal{R}^*L$  (see [7]).

For each  $\top_{\beta L} \neq I \in \beta L$ , the ideal  $M^I$  of  $\mathcal{R}L$  defined by

$$M^I = \{\varphi \in \mathcal{R}L : r_L(\text{coz}(\varphi)) \subseteq I\}$$

and  $M^{*I} = M^I \cap \mathcal{R}^*L$ . Also,

$$M^{*I} = \{\varphi \in \mathcal{R}^*L : \text{coz}(\varphi^\beta) \subseteq I\}.$$

We need the following propositions which are proved in [7].

**Proposition 5.2.** [7, Proposition 3.8] *Maximal ideals of  $\mathcal{R}^*L$  are precisely the ideals  $M^{*I}$ , for  $I \in \text{pt}(\beta L)$ . They are distinct for distinct  $I$ .*

**Proposition 5.3.** [7, Proposition 3.9] *For every  $I \in \text{pt}(\beta L)$ ,  $M^{*I}$  is fixed maximal ideal in  $\mathcal{R}^*L$  if and only if  $\bigvee I < \top$ .*

The following lemma plays an important role in this note.

**Lemma 5.4.** [10, Lemma 4.2] *For every  $p \in \text{pt}(L)$  and  $\varphi \in \mathcal{R}L$ ,  $\varphi[p] = 0$  if and only if  $\text{coz}(\varphi) \leq p$ .*

*Remark 5.5.* For every frame  $L$ , we put

$$L^* = \{I \in \text{pt}(\beta L) : \bigvee I = \top\}.$$

Also, for every  $A \subseteq \text{pt}(L)$  and  $\varphi \in \mathcal{R}L$ ,  $\varphi[A] = \{\varphi[p] : p \in A\}$ .

**Proposition 5.6.** *For every  $\varphi \in \mathcal{R}^*L$ , the following statements are equivalent:*

- (1)  $\varphi \in \bigcap_{I \in L^*} M^{*I}$ ;
- (2)  $\varphi^\beta[L^*] = \{0\}$ ;
- (3) For every  $0 < \varepsilon \in \mathbb{Q}$  and  $I \in L^*$ ,  $|\varphi^\beta[I]| < \varepsilon$ ;
- (4) For every  $n \in \mathbb{N}$ ,

$$\{I \in \text{pt}(\beta L) \mid |\varphi^\beta[I]| \geq \frac{1}{n}\} = \{I \in \text{pt}(\beta L) - L^* \mid |\varphi^\beta[I]| \geq \frac{1}{n}\}.$$

*Proof.* (1)  $\Leftrightarrow$  (2). By Lemma 5.4, we have

$$\begin{aligned} \varphi \in \bigcap_{I \in L^*} M^{*I} &\Leftrightarrow \forall I \in L^* (\text{coz}(\varphi^\beta) \subseteq I) \\ &\Leftrightarrow \forall I \in L^* (\varphi^\beta[I] = 0) \\ &\Leftrightarrow \varphi^\beta[L^*] = \{0\}. \end{aligned}$$

The rest is straightforward.  $\square$

**Theorem 5.7.** *The ring  $\mathcal{R}_\infty L$  is the intersection of all the free maximal ideals in  $\mathcal{R}^* L$ .*

*Proof.* Let  $\varphi \in \mathcal{R}_\infty L$  and  $I \in L^*$  such that  $\varphi \notin M^{*I}$ . Then

$$\bigvee_{n \in \mathbb{N}} \varphi^\beta((- , -\frac{1}{n}) \vee (\frac{1}{n}, -)) = \text{coz}(\varphi^\beta) \not\subseteq I.$$

So, there exists  $n_0 \in \mathbb{N}$  such that

$$\varphi^\beta((- , -\frac{1}{n_0}) \vee (\frac{1}{n_0}, -)) \not\subseteq I,$$

which implies that

$$\varphi^\beta((- , -\frac{1}{n_0}) \vee (\frac{1}{n_0}, -)) \vee I = \top_{\beta L}$$

and there exists  $a \in I$  and

$$x \in \varphi^\beta((- , -\frac{1}{n_0}) \vee (\frac{1}{n_0}, -))$$

such that  $x \vee a = \top$ . Since

$$x \leq \bigvee \varphi^\beta((- , -\frac{1}{n_0}) \vee (\frac{1}{n_0}, -)) = \varphi((- , -\frac{1}{n_0}) \vee (\frac{1}{n_0}, -)),$$

we conclude that

$$\varphi((- , -\frac{1}{n_0}) \vee (\frac{1}{n_0}, -)) \vee a = \top,$$

which implies

$$\varphi(-\frac{1}{n_0}, \frac{1}{n_0}) \leq (\varphi((- , -\frac{1}{n_0}) \vee (\frac{1}{n_0}, -)))^* \leq a.$$

It is clear that

$$A = \{x \vee a : x \in I\} \subseteq \uparrow \varphi(-\frac{1}{n_0}, \frac{1}{n_0})$$

and  $\bigvee A = \top$ . Since  $\uparrow \varphi(-\frac{1}{n_0}, \frac{1}{n_0})$  is compact frame, we conclude that there exist  $x_1, \dots, x_m \in I$  such that

$$\top = \bigvee_{i=1}^m (x_i \vee a) \in I,$$

which is a contradiction.

Conversely, let  $\varphi \in \bigcap_{I \in L^*} M^{*I}$ ,  $n \in \mathbb{N}$  and

$$\{a_\lambda\}_{\lambda \in \Lambda} \subseteq \uparrow \varphi(-\frac{1}{n}, \frac{1}{n})$$

such that  $\bigvee_{\lambda \in \Lambda} a_\lambda = \top$ . Suppose that for every  $\Lambda' \subseteq \Lambda$ , if  $\Lambda'$  is finite set, then  $\bigvee_{\lambda \in \Lambda'} a_\lambda \neq \top$ . Hence, there exists  $I \in L^*$  such that  $\{a_\lambda\}_{\lambda \in \Lambda} \subseteq I$ . By the statement (4) of Proposition 5.6, we have  $\varphi^\beta[I] = 0$ , so that  $\text{coz}(\varphi^\beta) \subseteq I$ , by Lemma 5.4. Since

$$\varphi\left(-\frac{1}{n}, \frac{1}{n}\right) \leq a_\lambda \in I,$$

we conclude that

$$\bigvee \varphi^\beta\left(-\frac{1}{n}, \frac{1}{n}\right) = \varphi\left(-\frac{1}{n}, \frac{1}{n}\right) \in I,$$

which follows that

$$\varphi^\beta\left(-\frac{1}{n}, \frac{1}{n}\right) \subseteq I.$$

Therefore,

$$L = \varphi^\beta\left(-\frac{1}{n}, \frac{1}{n}\right) \vee \text{coz}(\varphi^\beta) \subseteq I,$$

i.e.,  $L = I \in L^*$ , which is a contradiction.  $\square$

## 6. MAXIMAL IDEALS OF $\mathcal{R}_\infty L$

We turn our attention now to the fixed maximal ideals in the rings  $\mathcal{R}_\infty L$ .

**Lemma 6.1.** *Let  $\varphi \in \mathcal{R}L$ ,  $p \in \text{pt}(L)$  and  $n \in \mathbb{N}$ , then  $\varphi\left(\frac{-1}{n}, \frac{1}{n}\right) \leq p$  if and only if  $|\varphi[p]| \geq \frac{1}{n}$ .*

*Proof. Necessity.*

Let  $\varphi\left(\frac{-1}{n}, \frac{1}{n}\right) \leq p$  and  $|\varphi[p]| < \frac{1}{n}$ . If  $t = \varphi[p]$ , then, by Proposition 2.1,

$$\bigvee \{\varphi(-, r) \vee \varphi(s, -) : r, s \in \mathbb{Q}, r < t < s\} \leq p,$$

it follows that

$$\top = \varphi\left(\frac{-1}{n}, \frac{1}{n}\right) \vee \bigvee \{\varphi(-, r) \vee \varphi(s, -) : r, s \in \mathbb{Q}, r < t < s\} \leq p,$$

which is a contradiction.

*Sufficiency.* Let  $|\varphi[p]| \geq \frac{1}{n}$ . Then, by Proposition 2.1,

$$\varphi\left(\frac{-1}{n}, \frac{1}{n}\right) \leq \bigvee \{\varphi(-, r) \vee \varphi(s, -) | r, s \in \mathbb{Q}, r < \varphi[p] < s\} \leq p.$$

This completes the proof of the lemma.  $\square$

**Proposition 6.2.** *For every  $A \subseteq pt(L)$ , then  $\varphi[A] = 0$  for every  $\varphi \in \mathcal{R}_\infty L$ , if and only if for every  $\varphi \in \mathcal{R}_\infty L$  and  $n \in \mathbb{N}$ , if  $p \in A$ , then  $p \notin \uparrow \varphi(\frac{-1}{n}, \frac{1}{n})$ .*

*Proof. Necessity.* Let  $\varphi \in \mathcal{R}_\infty L$ ,  $p \in A$  and  $n \in \mathbb{N}$ . Suppose that  $p \in \uparrow \varphi(\frac{-1}{n}, \frac{1}{n})$ . Then, by Lemma 6.1,  $|\varphi[p]| \geq \frac{1}{n}$ . Hence,  $\varphi[p] \neq 0$ , which is a contradiction.

*Sufficiency.* Let  $\varphi \in \mathcal{R}_\infty L$  and  $p \in A$ . By Lemma 6.1,  $|\varphi[p]| < \frac{1}{n}$ , for every  $n \in \mathbb{N}$ . Hence  $\varphi[p] = 0$ .  $\square$

For each  $a \in L$  with  $a < \top$ , define the subset  $M_a$  of  $\mathcal{R}L$  by

$$M_a = \{\varphi \in \mathcal{R}L : \text{coz}(\varphi) \leq a\}$$

and  $M_a^* = M_a \cap \mathcal{R}^*L$ . Clearly,  $M_a$  is an ideal, and, in fact,  $M_a = M^{rL(a)}$ .

**Corollary 6.3.** *If  $p \in pt(L)$  then,  $\mathcal{R}_\infty L \subseteq M_p^*$  if and only if for every  $\varphi \in \mathcal{R}_\infty L$  and  $n \in \mathbb{N}$ ,  $p \notin \uparrow \varphi(\frac{-1}{n}, \frac{1}{n})$ .*

*Proof.* By Proposition 6.2, it is clear.  $\square$

For a proof of the following proposition, see [19, Corollary 3.6].

**Proposition 6.4.** *Let  $A$  be a commutative algebra over the rational numbers with unity. Let  $I$  be an ideal of  $A$ . Then an ideal  $D$  of  $I$  is a maximal ideal of  $I$  if and only if  $D = M \cap I$  for some maximal ideal  $M$  in  $A$ , with  $I \not\subseteq M$ .*

An ideal  $I$  in a subalgebra  $A$  of  $\mathcal{R}L$  is called strongly fixed ideal if  $\bigcap_{\varphi \in I} Z(\varphi) \neq \emptyset$ , otherwise,  $I$  is said to be strongly free ideal.

For a proof of the following proposition, see [7, Proposition 3.3] or [10, Proposition 4.8, Corollary 4.9].

**Proposition 6.5.** *The fixed maximal ideals of  $\mathcal{R}L$  ( $\mathcal{R}^*L$ ) are precisely the ideals  $M_p$  ( $M_p^*$ ) for  $p \in Pt(L)$ . They are distinct for distinct points.*

**Proposition 6.6.** *If  $L$  is a completely regular frame, then every maximal ideal of  $\mathcal{R}_\infty L$  is strongly fixed ideal. In fact,  $M$  is a maximal ideal of  $\mathcal{R}_\infty L$  if and only if there exists  $p \in pt(L)$  such that*

- (1)  $M = M_p^* \cap \mathcal{R}_\infty L$ , and
- (2)  $p \in \uparrow \varphi(\frac{-1}{n}, \frac{1}{n})$ , for some  $\varphi \in \mathcal{R}_\infty L$  and  $n \in \mathbb{N}$ .

*Proof.* Let  $M$  be a maximal ideal of  $\mathcal{R}_\infty L$ , then by Propositions 5.2 and 6.4, there exists  $I \in pt(\beta L)$  such that  $M = M^{*I} \cap \mathcal{R}_\infty L$ , with  $\mathcal{R}_\infty L \not\subseteq M^{*I}$ . By Theorem 5.7,  $M^{*I}$  is a fixed maximal ideal of  $\mathcal{R}^*L$ .

Then, there exists  $p \in pt(L)$  such that  $M^{*I} = M_p^*$ , by Proposition 6.5. Therefore, we have

- (1)  $M = M_p^* \cap \mathcal{R}_\infty L$ , and
- (2)  $p \in \uparrow \varphi(\frac{-1}{n}, \frac{1}{n})$ , for some  $\varphi \in \mathcal{R}_\infty L$  and  $n \in \mathbb{N}$ , by Corollary 6.3.

Conversely, by Corollary 6.3 and Propositions 6.4 and 6.5, it is clear that  $M$  is a maximal ideal of  $\mathcal{R}_\infty L$ .  $\square$

#### ACKNOWLEDGMENTS

The authors thank the anonymous referees for their valuable comments and suggestions for improving the paper.

#### REFERENCES

- [1] S. K. Acharyya, K. C. Chattopadhyay and D. P. Ghosh, On a class of subalgebras of  $C(X)$  and the intersection of their free maximal ideals, *Proc. Amer. Math. Soc.* **125**(2) (1997), 611–615.
- [2] S. K. Acharyya and D. De, An interesting class of ideals in subalgebras of  $C(X)$  containing  $C^*(X)$ , *Comment. Math. Univ. Carolin.* **48**(2) (2007), 273–280.
- [3] S. Afrooz and M. Namdari,  $C_\infty(X)$  and related ideals, *Real Anal. Exchange* **36**(1) (2010/2011), 45–54.
- [4] F. Azarpanah and R. Soundararajan, When the family of functions vanishing at infinity is an ideal of  $C_\infty(X)$ , *Rocky Mountain J. Math.* **31**(4) (2001), 1133–1140.
- [5] B. Banaschewski, The real numbers in pointfree topology, *Textos de Matematica (Series B)* **12** (1997), 1–96.
- [6] T. Dube, On the ideal of functions with compact support in pointfree function rings, *Acta Math. Hungar.* **129**(3) (2010), 205–226.
- [7] T. Dube, Extending and contracting maximal ideals in the function rings of pointfree topology, *Bull. Math. Soc. Sci. Math. Roumanie* **55**(4) (2012), 365–374.
- [8] M. M. Ebrahimi and A. Karimi Feizabadi, Prime representation of real Riesz maps, *Algebra Universalis* **54** (2005), 291–299.
- [9] M. M. Ebrahimi and M. Mahmoudi, *Frame*, Technical Report, Shahid Beheshti University, 1996.
- [10] A. A. Estaji, A. Karimi Feizabadi and M. Abedi, Strongly fixed ideals in  $C(L)$  and compact frames, *Arch. Math. (Brno)* **51** (2015), 1–12.
- [11] A. A. Estaji, A. Karimi Feizabadi and M. Abedi, Zero set in pointfree topology and strongly  $z$ -ideals, *Bull. Iranian Math. Soc.* **41**, No.5 (2015), 1071–1084.
- [12] A. A. Estaji, A. Karimi Feizabadi and M. Abedi, Intersection of essential ideals in the ring of real-valued continuous functions on a frame, *Journal of Algebraic Systems* **5**, No.2 (2017), 149–161.
- [13] A. A. Estaji, A. Karimi Feizabadi and M. Robat Sarpoushi,  $z_c$ -ideals and prime ideals in the ring  $\mathcal{R}_c L$ , *Filomat*, in press.
- [14] L. Gillman and M. Jerison, *Rings of continuous functions*, Springer-Verlag, 1976.

- [15] A. Karimi Feizabadi, A. A. Estaji and M. Abedi, On minimal ideals in the ring of real-valued continuous functions on a frame, *Arch. Math. (Brno)* Accepted.
- [16] C. W. Kohls, Ideals in rings of continuous functions, *Fund. Math.* **45** (1957), 28–50.
- [17] J. Picado and A. Pultr, *Frames and locales: Topology without points*, Frontiers in Mathematics, Springer Basel, 2012.
- [18] A. Rezaei Aliabad, F. Azarpanah and M. Namdari,, Rings of continuous functions vanishing at infinity, *Comment. Math. Univ. Carolin.* **45**(3) (2004), 519–533.
- [19] D. Rudd, On isomorphisms between ideals in rings of continuous functions, *Trans. Amer. Math. Soc.* **159** (1971), 335–353.
- [20] A. Taherifar, Some generalizations and unifications of  $C_K(X)$ ,  $C_\psi(X)$  and  $C_\infty(X)$ , *Quaest. Math.* **38** (2015), 793–804.

**Ali Akbar Estaji**

Faculty of Mathematics and Computer Sciences, Hakim Sabzevari University, Sabzevar, Iran.

Email: [aaestaji@hsu.ac.ir](mailto:aaestaji@hsu.ac.ir) and [aaestaji@gmail.com](mailto:aaestaji@gmail.com)

**Ahmad Mahmoudi Darghadam**

Faculty of Mathematics and Computer Sciences, Hakim Sabzevari University, Sabzevar, Iran.

Email: [m.darghadam@yahoo.com](mailto:m.darghadam@yahoo.com)

On maximal ideals of  $\mathcal{R}_\infty L$

Ali Akbar Estaji and Ahmad Mahmoudi Darghadam

ایده‌آل‌های ماکسیمال  $\mathcal{R}_\infty L$

علی اکبر استاجی و احمد محمودی درقدم  
دانشکده ریاضی و علوم کامپیوتر دانشگاه حکیم سبزواری، ایران، سبزواری

فرض کنید  $L$  قاب کاملاً منظم و  $\mathcal{R}L$  حلقه توابع پیوسته حقیقی مقدار روی قاب  $L$  باشد. قرار می‌دهیم

$$\mathcal{R}_\infty L = \{ \varphi \in \mathcal{R}L : \varphi \uparrow \left( \frac{-1}{n}, \frac{1}{n} \right), n \in \mathbb{N} \text{ هر برای} \}.$$

فرض کنید  $C_\infty(X)$  گردایه‌ی تمام عناصر  $f \in C(X)$  به قسمی باشند که برای هر  $n \in \mathbb{N}$ ،  $\{x \in X : |f(x)| \geq \frac{1}{n}\}$  فشرده است. کلز نشان داد که  $C_\infty(X)$  برابر با اشتراک تمام ایده‌آل‌های ماکسیمال آزاد  $C^*(X)$  است. در این مقاله می‌خواهیم این نتیجه را به حلقه توابع پیوسته حقیقی مقدار روی قاب گسترش دهیم و نشان می‌دهیم  $\mathcal{R}_\infty L$  دقیقاً برابر با اشتراک تمام ایده‌آل‌های ماکسیمال آزاد  $\mathcal{R}^*L$  است. با استفاده از این نتیجه، ایده‌آل‌های ماکسیمال  $\mathcal{R}_\infty L$  را شناسایی می‌کنیم.

کلمات کلیدی: قاب، فشرده، ایده‌آل ماکسیمال، حلقه توابع پیوسته حقیقی مقدار.