

COMPUTING THE PRODUCTS OF CONJUGACY CLASSES FOR SPECIFIC FINITE GROUPS

MARYAM JALALI-RAD AND ALI REZA ASHRAFI*

ABSTRACT. Suppose G is a finite group, A and B are conjugacy classes of G , and $\eta(AB)$ denotes the number of conjugacy classes contained in AB . The set of all $\eta(AB)$, such that A, B run over conjugacy classes of G , is denoted by $\eta(G)$. The aim of this paper is to compute $\eta(G)$, for $G \in \{D_{2n}, T_{4n}, U_{6n}, V_{8n}, SD_{8n}\}$ or G is a decomposable group of order $2pq$, a group of order $4p$ or p^3 , where p and q are primes.

1. INTRODUCTION

Throughout this paper, all groups are assumed to be finite. If G is such a group, and A and B are conjugacy classes of G , then it is an elementary fact that AB is a G -invariant subset. Thus, AB can be written as a union of conjugacy classes of G . The number of distinct conjugacy classes of G contained in AB is denoted by $\eta(AB)$. The set of all $\eta(AB)$, such that A, B run over conjugacy classes of G , is denoted by $\eta(G)$.

The most important works on the problem of computing the number of G -conjugacy classes in the product of conjugacy classes were carried out by Adan–Bante. Here, we report some of her interesting results in this topic. Suppose $SL(2, q)$ is the group of 2×2 matrices, with determinant one over a finite field of order q . Adan–Bante and Harris [3] proved that if q is even, then the product of any two non-central conjugacy classes of $SL(2, q)$ is a union of at least $q - 1$ distinct

MSC(2010): Primary: 20E45; Secondary: 20C15

Keywords: Conjugacy class, G -invariant subset, p -group.

Received: 30 November 2014, Revised: 27 April 2015.

*Corresponding author .

conjugacy classes of $SL(2, q)$; and if $q > 3$ is odd, then the product of any two non-central conjugacy classes of $SL(2, q)$ is the union of at least $\frac{q+3}{2}$ distinct conjugacy classes of $SL(2, q)$. Adan–Bante [1] proved that, for any finite supersolvable group G , and any conjugacy class A of G , $dl(\frac{G}{C_G(A)}) \leq 2\eta(AA^{-1}) - 1$, where $C_G(A)$ denotes the centralizer of A in G , and $dl(H)$ is the derived length of a group H . In [2], she also proved that if p is an odd prime number, G is a finite p -group, and a^G and b^G are the conjugacy classes of G of size p ; then either $a^G b^G = (ab)^G$ or $a^G b^G$ is a union of at least $\frac{p+1}{2}$ distinct conjugacy classes. If G is nilpotent, and a^G is again a conjugacy class of G of size p , then either $a^G a^G = (a^2)^G$ or $a^G a^G$ is a union of exactly $\frac{p+1}{2}$ distinct conjugacy classes of G of size p .

Darafsheh and Robati [6] continued the works of Adan–Bante and proved that if $[a, G] = \{[a, x] \mid x \in G\}$, and $[a, G]$ be a subset of $Z(G)$, then we have:

- i. $\eta(a^G b^G) = |a^G| |b^G| / |[a, G] \cap (b^{-1})^G b^G| |(ab)^G|$;
- ii. If $a^G b^G \cap Z(G) \neq \emptyset$, then $\eta(a^G b^G) = |a^G|$;
- iii. If $|a^G|$ is an odd number, then $\eta(a^G a^G) = 1$;
- iv. If $|a^G|$ is an even number, then $\eta(a^G a^G) = 2^n$, where n is the number of cyclic direct factors in the decomposition of the Sylow 2-subgroup of $[a, G]$.

We encourage the interested readers to consult also the papers by Arad and his co-authors [4, 5], and references therein for more information on this topic. Our notation is standard, and can be taken from [9, 10].

2. MAIN RESULTS

The aim of this section is to compute $\eta(G)$, where

$$G \in \{D_{2n}, V_{8n}, T_{4n}, U_{6n}, SD_{8n}\}$$

or G is a group of orders $2pq, 4p, p^3$, such that p and q are prime numbers. The case of $|G| = 2pq$ and G that is indecomposable, is retained as an open question. The semi-dihedral group SD_{8n} , dicyclic group T_{4n} , and the groups U_{6n} and V_{8n} have the following presentations, respectively:

$$\begin{aligned} SD_{8n} &= \langle a, b \mid a^{4n} = b^2 = e, bab = a^{2n-1} \rangle, \\ T_{4n} &= \langle a, b \mid a^{2n} = 1, a^n = b^2, b^{-1}ab = a^{-1} \rangle, \\ U_{6n} &= \langle a, b \mid a^{2n} = b^3 = e, bab = a \rangle, \\ V_{8n} &= \langle a, b \mid a^{2n} = b^4 = e, aba = b^{-1}, ab^{-1}a = b \rangle. \end{aligned}$$

It is easy to see the dicyclic group T_{4n} has the order $4n$, and the cyclic subgroup $\langle a \rangle$ of T_{4n} has the index 2 [10]. The conjugacy classes of U_{6n} and V_{8n} (n is odd), computed in the famous book of James and Liebeck [10]. The groups V_{8n} (n is even), and SD_{8n} have the order $8n$, and their conjugacy classes have been computed in [7, 8], respectively.

The following simple lemma is crucial throughout this paper:

Lemma 2.1. *Suppose G is a finite group, and A and B are conjugacy classes of G . Then,*

- (1) $\eta(AB) = \eta(BA)$,
- (2) *If A is central, then $\eta(AB) = 1$,*
- (3) *If $|A| = |B| = 2$, then $\eta(AB) = 1, 2$ [2, Proposition 2.7],*
- (4) $\eta(AB) \leq |A|$ [6, Lemma 3.1].

Proposition 2.2.

$$\eta(D_{2n}) = \begin{cases} \{1, 2, \frac{n+1}{2}\} & 2 \nmid n \\ \{1, 2, \frac{n}{4}, \frac{n}{4} + 1\} & n \equiv 0 \pmod{4} \\ \{1, 2, \frac{n+2}{4}\} & n \equiv 2 \pmod{4} \end{cases} .$$

Proof. The dihedral group D_{2n} can be presented by

$$D_{2n} = \langle a, b \mid a^n = b^2 = e, b^{-1}ab = a^{-1} \rangle.$$

We first assume that n is odd. Then the conjugacy classes of D_{2n} are $\{e\}, \{a^r, a^{-r}\}, 1 \leq r \leq \frac{n-1}{2}$ or $\{a^s b; 0 \leq s \leq n-1\}$. Thus the products of non-identity conjugacy classes are:

- $\{a^r, a^{-r}\} \cdot \{a^s, a^{-s}\} = \{a^{r+s}, a^{-(r+s)}\} \cup \{a^{r-s}, a^{-(r-s)}\}$,
- $\{a^r, a^{-r}\} \cdot \{a^s b; 0 \leq s \leq n-1\} = \{a^{r+s} b, a^{s-r} b; 0 \leq s \leq n-1\} = \{a^s b; 0 \leq s \leq n-1\}$,
- $\{a^s b; 0 \leq s \leq n-1\} \cdot \{a^r b; 0 \leq r \leq n-1\} = \{a^s b a^r b; 0 \leq r, s \leq n-1\} = \bigcup_{r=0}^{\frac{n-1}{2}} \{a^r, a^{-r}\}$.

Hence, $\eta(D_{2n}) = \{1, 2, \frac{n+1}{2}\}$. Next, assume that $n = 2m$. The conjugacy classes of D_{2n} are $\{e\}, \{a^m\}, \{a^r, a^{-r}\}, 1 \leq r \leq m-1, \{a^s b; 0 \leq s \leq 2(n-1), 2 \mid s\}, \{a^s b; 0 \leq s \leq 2(n-1), 2 \nmid s\}$. Suppose $0 \leq r, l \leq m-1, F_1 = \{0 \leq s \leq 2(n-1), 2 \mid s\}$, and $F_2 = \{0 \leq s \leq 2(n-1), 2 \nmid s\}$. The products of non-identity conjugacy classes are as follows:

$$\begin{aligned}
\{a^r, a^{-r}\} \cdot \{a^l, a^{-l}\} &= \{a^{r-l}, a^{l-r}\} \cup \{a^{r+l}, a^{-(l+r)}\}, \\
\{a^r, a^{-r}\} \cdot \{a^s b; s \in F_1\} &= \begin{cases} \{a^s b; s \in F_1\} & 2 \mid r \\ \{a^s b; s \in F_2\} & 2 \nmid r \end{cases}, \\
\{a^r, a^{-r}\} \cdot \{a^s b; s \in F_2\} &= \begin{cases} \{a^s b; s \in F_1\} & 2 \nmid r \\ \{a^s b; s \in F_2\} & 2 \mid r \end{cases}, \\
\{a^s b; s \in F_1\} \cdot \{a^r b; r \in F_1\} &= \{e\} \cup \begin{cases} \bigcup_{r=1}^{\frac{n-2}{4}} \{a^{2r}, a^{-2r}\} & n \equiv 2 \pmod{4} \\ \bigcup_{r=1}^{\frac{n}{4}} \{a^{2r}, a^{-2r}\} & n \equiv 0 \pmod{4} \end{cases}, \\
\{a^s b; s \in F_1\} \cdot \{a^r b; r \in F_2\} &= \{e\} \cup \begin{cases} \bigcup_{r=1}^{\frac{n-2}{4}} \{a^{2r}, a^{-2r}\} & n \equiv 0 \pmod{4} \\ \bigcup_{r=1}^{\frac{n}{4}} \{a^{2r}, a^{-2r}\} & n \equiv 2 \pmod{4} \end{cases}, \\
\{a^s b; s \in F_2\} \cdot \{a^r b; r \in F_2\} &= \{e\} \cup \begin{cases} \bigcup_{r=0}^{\frac{n-6}{4}} (a^{2r+1})^{D_{2n}} & n \equiv 2 \pmod{4} \\ \bigcup_{r=0}^{\frac{n}{4}} (a^{2r+1})^{D_{2n}} & n \equiv 0 \pmod{4} \end{cases}.
\end{aligned}$$

This completes the proof. \square

Proposition 2.3.

$$\eta(V_{8n}) = \begin{cases} \{1, 2, \frac{n}{2}, \frac{n}{2} + 1\} & n \text{ is even} \\ \{1, 2, n, n + 1\} & n \text{ is odd} \end{cases}.$$

Proof. By Lemma 2.1 (1, 2), it is enough to compute $\eta(AB)$, where A and B are the non-central conjugacy classes of V_{8n} . Our main proof considers two separate cases, in which n is odd or even.

We first assume that n is odd. Then by [10], the conjugacy classes of V_{8n} are as follows:

$$\begin{aligned}
&\{e\}, \{b^2\}, \{a^{2r+1}, a^{-(2r+1)}b^2\}, 0 \leq r \leq \frac{n-1}{2}, \{a^{2s}, a^{-2s}\}, \{a^{2s}b^2, a^{-2s}b^2\}, \\
&1 \leq s \leq \frac{n-1}{2}, \{a^j b^k; k = 1, 3 \ \& \ 2 \mid j\} \text{ and } \{a^j b^k; k = 1, 3 \ \& \ 2 \nmid j\}.
\end{aligned}$$

Before starting our calculations, we notice that if A and B are two conjugacy classes of length 2, then by Lemma 2.1 (3), $\eta(AB) = 2$. Thus, it is enough to consider the cases where $(|A|, |B|) \neq (2, 2)$.

- $(a^{2s}, a^{-2s}) \cdot \{a^j b^k; k = 1, 3\} = \{a^{j+2s} b^k; k = 1, 3\} \cup \{a^{j-2s} b^k; k = 1, 3\}$,
- $\{a^{2r+1}, a^{-(2r+1)}b^2\} \cdot \{a^j b^k; k = 1, 3 \ \& \ 2 \nmid j\} = \{a^j b^k; k = 1, 3 \ \& \ 2 \mid j\} \cup \{a^j b^k; k = 1, 3 \ \& \ 2 \nmid j\}$,
- $\{a^{2r+1}, a^{-(2r+1)}b^2\} \cdot \{a^j b^k; k = 1, 3 \ \& \ 2 \mid j\} = \{a^j b^k; k = 1, 3 \ \& \ 2 \nmid j\}$,
- $\{a^{2s}b^2, a^{-2s}b^2\} \cdot \{a^j b^k; k = 1, 3 \ \& \ 2 \mid j\} = \{a^j b^k; k = 1, 3 \ \& \ 2 \mid j\}$,
- $(b)^{V_{8n}} \cdot (b)^{V_{8n}} = \bigcup_{s=0}^{\frac{n-1}{2}} \{a^{2s}, a^{-2s}\} \cup \bigcup_{s=0}^{\frac{n-1}{2}} \{a^{2s}b^2, a^{-2s}b^2\}$,
- $(ab)^{V_{8n}} \cdot (ab)^{V_{8n}} = \bigcup_{s=0}^{\frac{n-1}{2}} \{a^{2s}, a^{-2s}\} \cup \bigcup_{s=0}^{\frac{n-1}{2}} \{a^{2s}b^2, a^{-2s}b^2\}$,
- $(b)^{V_{8n}} \cdot (ab)^{V_{8n}} = \bigcup_{r=0}^{\frac{n-1}{2}} \{a^{2r+1}, a^{-(2r+1)}b^2\}$.

Next, we assume that $n = 2l$ is even. Then, by [6], the conjugacy classes of V_{8n} are $\{e\}, \{b^2\}, \{a^n\}, \{a^n b^2\}, \{a^{2k+1}b^{(-1)^{k+1}}; 0 \leq k \leq n-1\}, \{a^{2r+1}, a^{-(2r+1)}b^2\}, 0 \leq r \leq n-1, \{a^{2s}, a^{-2s}\}, \{a^{2s}b^2, a^{-2s}b^2\}$,

$1 \leq s \leq \frac{n}{2} - 1$, $\{a^{2k}b^{(-1)^k}; 0 \leq k \leq n-1\}$, $\{a^{2k}b^{(-1)^{k+1}}; 0 \leq k \leq n-1\}$, $\{a^{2k+1}b^{(-1)^k}; 0 \leq k \leq n-1\}$. Suppose $0 \leq k \leq n-1$, and $0 \leq r, s \leq \frac{n}{2} - 1$. Then the product of non-central conjugacy classes are as follows:

$$\bullet \{a^{2s}, a^{-2s}\} \cdot \{a^{2r}, a^{-2r}\} = \{a^{2(r+s)}, a^{-2(r+s)}\} \cup \{a^{2(r-s)}, a^{-2(r-s)}\}.$$

\bullet Suppose:

$$F = \{b^2\} \cup \{a^n b^2\} \cup_{r=1, 2 \nmid r}^{\frac{n}{2}-1} \{a^{2s}, a^{-2s}\} \cup_{r=1, 2 \nmid r}^{\frac{n}{2}-1} \{a^{2s}b^2, a^{-2s}b^2\}.$$

Then $\{a^{2k}b^{(-1)^k}; 0 \leq k \leq n-1\} \cdot \{a^{2k}b^{(-1)^k}; 0 \leq k \leq n-1\}$ can be simplified as follows:

$$F \cup \begin{cases} \{a^n b^2\} & n \equiv 0 \pmod{4} \\ \{a^n\} & n \equiv 2 \pmod{4} \end{cases}.$$

Therefore, $\eta(b^{V_{8n}} \cdot b^{V_{8n}}) = \frac{n}{2} + 1$.

\bullet In a similar argument as above, we have:

$$\begin{aligned} \eta((b)^{V_{8n}} \cdot (b^{-1})^{V_{8n}}) &= \eta((ab^{-1})^{V_{8n}} \cdot (ab^{-1})^{V_{8n}}) \\ &= \eta((ab^{-1})^{V_{8n}} \cdot (ab)^{V_{8n}}) \\ &= \eta((b^{-1})^{V_{8n}} \cdot (b^{-1})^{V_{8n}}) \\ &= \eta((b^{-1})^{V_{8n}} \cdot (ab)^{V_{8n}}) \\ &= \eta((ab)^{V_{8n}} \cdot (ab)^{V_{8n}}) = \frac{n}{2} + 1. \end{aligned}$$

\bullet In the following case, it can be proved that $\eta((ab^{-1})^{V_{8n}} \cdot b^{V_{8n}}) = \frac{n}{2}$.

$$(ab^{-1})^{V_{8n}} \cdot (b^{-1})^{V_{8n}} = \bigcup_{r=1, 2 \nmid r}^{n-1} \{a^{2r+1}, a^{-(2r+1)}b^2\}.$$

\bullet For the following product of conjugacy classes, we have:

$$\eta((ab^{-1})^{V_{8n}} \cdot (b)^{V_{8n}}) = \eta((b)^{V_{8n}} \cdot (ab)^{V_{8n}}) = \frac{n}{2}.$$

\bullet

$$(a^{2r+1})^{V_{8n}} \cdot (b^{-1})^{V_{8n}} = \begin{cases} (ab^{-1})^{V_{8n}} & r \equiv 1 \pmod{4} \\ (b^{-1})^{V_{8n}} & r \equiv 3 \pmod{4} \end{cases}.$$

\bullet

$$(a^{2r+1})^{V_{8n}} \cdot (ab)^{V_{8n}} = \begin{cases} (b^{-1})^{V_{8n}} & r \equiv 1 \pmod{4} \\ (b)^{V_{8n}} & r \equiv 3 \pmod{4} \end{cases}.$$

\bullet

$$(a^{2r+1})^{V_{8n}} \cdot (ab^{-1})^{V_{8n}} = \begin{cases} (b)^{V_{8n}} & r \equiv 1 \pmod{4} \\ (ab)^{V_{8n}} & r \equiv 3 \pmod{4} \end{cases}.$$

•

$$(a^{2r+1})^{V_{8n}} \cdot (b)^{V_{8n}} = \begin{cases} (ab)^{V_{8n}} & r \equiv 1 \pmod{4} \\ (ab^{-1})^{V_{8n}} & r \equiv 1 \pmod{4} \end{cases}.$$

The product of Conjugacy classes of length two by another conjugacy class of two given types is again a conjugacy class. This completes the proof. \square

Proposition 2.4.

$$\eta(T_{4n}) = \begin{cases} \{1, 2, \frac{n}{2}, \frac{n}{2} + 1\} & n \text{ is even} \\ \{1, 2, \frac{n+1}{2}\} & n \text{ is odd} \end{cases}.$$

Proof. By [10, p. 420], the conjugacy classes of T_{4n} are $\{e\}$, $\{a^n\}$, $\{a^r, a^{-r}\}$, $1 \leq r \leq n-1$, $\{a^{2j}b, 0 \leq j \leq n-1\}$, $\{a^{2j+1}b, 0 \leq j \leq n-1\}$. On the other hand, the product of conjugacy classes can be computed, as follows:

- $\{a^r, a^{-r}\} \cdot \{a^s, a^{-s}\} = \{a^{r+s}, a^{-(r+s)}\} \cup \{a^{r-s}, a^{-(r-s)}\}$.
- Since $(a^r)^{T_{4n}} \cdot (b)^{T_{4n}} = \{a^{r+2j}b, a^{-r+2j}b; 0 \leq j \leq n-1\}$, the product is $(b)^{T_{4n}}$, when r is even. If r is odd, then the product will be $(ab)^{T_{4n}}$.
- We know that $(a^r)^{T_{4n}} \cdot (ab)^{T_{4n}} = \{a^{r+2j+1}b, a^{-r+2j+1}b; 0 \leq j \leq n-1\}$. If r is even, then the product is $(ab)^{T_{4n}}$, and if r is odd, then the product will be $(b)^{T_{4n}}$.

•

$$(b)^{T_{4n}} \cdot (b)^{T_{4n}} = \begin{cases} \bigcup_{r=0,2|r}^n \{a^r, a^{-r}\} & 2 \mid n \\ \bigcup_{r=1,2 \nmid r}^n \{a^r, a^{-r}\} & 2 \nmid n \end{cases}.$$

•

$$(b)^{T_{4n}} \cdot (ab)^{T_{4n}} = \begin{cases} \bigcup_{r=1,2 \nmid r}^{n-1} \{a^r, a^{-r}\} & 2 \mid n \\ \bigcup_{r=0,2 \nmid r}^{n-1} \{a^r, a^{-r}\} & 2 \nmid n \end{cases}.$$

•

$$(ab)^{T_{4n}} \cdot (ab)^{T_{4n}} = \begin{cases} \bigcup_{r=0,2|r}^n \{a^r, a^{-r}\} & 2 \mid n \\ \bigcup_{r=1,2 \nmid r}^n \{a^r, a^{-r}\} & 2 \nmid n \end{cases}.$$

This completes the proof. \square

Example 2.5. Suppose G is a non-abelian group of order $4p$; p is prime. By an easy calculation, one can see that $\eta(D_8) = \eta(Q_8) = \eta(D_{12}) = \eta(Z_3 : Z_4) = \eta(A_4) = \{1, 2\}$, where $Z_3 : Z_4$ is a non-abelian group of order 12 different from A_4 and D_{12} . Thus, it is enough to consider that case that $p > 3$. Our proof considers two cases Thus $4 \mid p-1$ or $4 \nmid p-1$.

Case 1. $4|p-1$. If $4|p-1$, then up to isomorphism, there are three groups of order $4p$. These are D_{4p} , T_{4p} , and F_{4p} , where F_{4p} can be presented by $F_{4p} = \langle a, b | a^p = b^4 = 1, b^{-1}ab = a^\lambda \rangle$, and $\lambda^2 \equiv -1 \pmod{p}$. By Propositions 2 and 4, $\eta(D_{4p}) = \eta(T_{4p}) = \{1, 2, \frac{p+1}{2}\}$, and $\eta(F_{4p}) = \{1, 3, 4, \frac{p+3}{4}\}$.

Case 2. $4 \nmid p-1$. In this case, there are up to isomorphism two groups of order $4p$. These are D_{4p} and T_{4p} . As in Case 1, $\eta(D_{4p}) = \eta(T_{4p}) = \{1, 2, \frac{p+1}{2}\}$, as desired.

Therefore,

$$\eta(G) \in \begin{cases} \{\{1, 2, \frac{p+1}{2}\}\} & p > 3, 4 \nmid p-1 \\ \{\{1, 2, \frac{p+1}{2}\}, \{1, 3, 4, \frac{p+3}{4}\}\} & p > 3, 4 | p-1 \\ \{\{1, 2\}\} & p = 3 \end{cases} .$$

Proposition 2.6. $\eta(U_{6n}) = \{1, 2\}$.

Proof. By [10], the conjugacy classes of U_{6n} are $\{e\}, \{a^{2r}\}, \{a^{2r}b, a^{2r}b^2\}, \{a^{2r+1}, a^{2r+1}b, a^{2r+1}b^2\}, 0 \leq r \leq n-1$. On the other hand, by Lemma 2.1 (4), $\eta((a^{2r}b)^{U_{6n}} \cdot (a^{2s+1})^{U_{6n}}) \leq 2$. But, $(a^{2r+1})^{U_{6n}} \cdot (a^{2s+1})^{U_{6n}} = \{a^{2(r+s+1)}\} \cup \{a^{2(r+s+1)}b, a^{2(r+s+1)}b^2\}$. Thus, $\eta(U_{6n}) = \{1, 2\}$, as desired. \square

Hormozi and Rodtes [8, Definition 2.1], defined $C^{even} = C_1 \cup C_2^{even} \cup C_3^{even}$ and $C^{odd} = C_1 \cup C_2^{odd} \cup C_3^{odd}$, where $C_1 = \{0, 2, 4, \dots, 2n\}$, $C_2^{even} = \{1, 3, 5, \dots, n-1\}$, $C_3^{even} = \{2n+1, 2n+3, 2n+5, \dots, 3n-1\}$, $C_2^{odd} = \{1, 3, 5, \dots, n\}$, $C_3^{odd} = \{2n+1, 2n+3, 2n+5, \dots, 3n\}$, $C_{even}^\dagger = C_1 \setminus \{0, 2n\}$ and $C_{odd}^\dagger = C_2^{even} \cup C_3^{even}$. Moreover, $C_*^{even} = C^{even} \setminus \{0, 2n\}$ and $C_*^{odd} = C^{odd} \setminus \{0, n, 2n, 3n\}$.

Proposition 2.7.

$$\eta(SD_{8n}) = \begin{cases} \{1, 2, n, n+1\} & n \text{ is even} \\ \{1, 2, \frac{n+1}{2}\} & n \text{ is odd} \end{cases} .$$

Proof. By [8, Proposition 2.2], the conjugacy classes of SD_{8n} ; $n \geq 2$, can be computed in two separate cases, where n is odd or even. If n is even, then there are $2n+3$ conjugacy classes as: $\{e\}, \{a^{2n}\}, \{a^r, a^{(2n-1)r}\}; r \in C_*^{even}, \{ba^{2t} | t = 0, 1, 2, \dots, 2n-1\}$ and $\{ba^{2t+1} | t = 0, 1, 2, \dots, 2n-1\}$. If n is odd, then there are $2n+6$ conjugacy classes as $\{e\}, \{a^n\}, \{a^{2n}\}, \{a^{3n}\}, \{a^r, a^{(2n-1)r}\}; r \in C_*^{odd}, \{ba^{4t} | t = 0, 1, 2, \dots, n-1\}, \{ba^{4t+1} | t = 0, 1, 2, \dots, n-1\}, \{ba^{4t+2} | t = 0, 1, 2, \dots, n-1\}$ and $\{ba^{4t+3} | t = 0, 1, 2, \dots, n-1\}$. On the other hand by [8], we have:

- (1) $(2n-1)r \equiv (4n-r) \pmod{4n}$, if r is even,
- (2) $(2n-1)r \equiv (2nr) \pmod{4n}$, if r is odd,

(3) $(2n - 1)(2n + k) \equiv (4n - k) \pmod{4n}$, if k is odd.

If n is even, then the product of conjugacy classes, of SD_{8n} are as follows:

- $\{a^r, a^{(2n-1)r} \& 2|r\} \cdot \{a^s, a^{(2n-1)s} \& 2|s\} = \{a^r, a^{(2n-1)r} \& 2 \nmid r\} \cdot \{a^s, a^{(2n-1)s} \& 2 \nmid s\} = (a^{r+s})^{SD_{8n}} \cup (a^{r+(2n-1)s})^{SD_{8n}}$,
- $\{a^r, a^{(2n-1)r} \& 2|r\} \cdot (b)^{SD_{8n}} = \{a^r, a^{(2n-1)r} \& 2 \nmid r\} \cdot (ba)^{SD_{8n}} = (b)^{SD_{8n}}$,
- $\{a^r, a^{(2n-1)r} \& 2|r\} \cdot (ba)^{SD_{8n}} = \{a^r, a^{(2n-1)r} \& 2 \nmid r\} \cdot (b)^{SD_{8n}} = (ba)^{SD_{8n}}$,
- $(b)^{SD_{8n}} \cdot (b)^{SD_{8n}} = (ba)^{SD_{8n}} \cdot (ba)^{SD_{8n}} = \bigcup_{r \in C_1} \{a^r, a^{(2n-1)r}\}$,
- $(b)^{SD_{8n}} \cdot (ba)^{SD_{8n}} = \bigcup_{r \in C_2^{even} \cup C_3^{even}} \{a^r, a^{(2n-1)r}\}$.

If n is odd, then the products of conjugacy classes of SD_{8n} are as follows:

- $(a^r)^{SD_{8n}} \cdot (a^s)^{SD_{8n}} = \{a^{r+s}, a^{(2n-1)(r+s)}\} \cup \{a^{r+(2n-1)s}, a^{s+(2n-1)r}\}$,
- $(a^r)^{SD_{8n}} \cdot (ba^i)^{SD_{8n}} = (ba^j)^{SD_{8n}}$, where $j = i - r$, $i + 2n - r$, when r is even or odd, respectively.
- $(b)^{SD_{8n}} \cdot (b)^{SD_{8n}} = (ba^2)^{SD_{8n}} \cdot (ba^2)^{SD_{8n}} = (ba)^{SD_{8n}} \cdot (ba^3)^{SD_{8n}} = \bigcup_{r \in C_1, r \equiv 0 \pmod{4}} \{a^r, a^{(2n-1)r}\}$,
- $(b)^{SD_{8n}} \cdot (ba)^{SD_{8n}} = \bigcup_{r \in C_2^{odd} \cup C_3^{odd}, r \equiv 1 \pmod{4}} \{a^r, a^{(2n-1)r}\}$,
- $(ba^2)^{SD_{8n}} \cdot (ba^3)^{SD_{8n}} = \bigcup_{r \in C_2^{odd} \cup C_3^{odd}, r \equiv 1 \pmod{4}} \{a^r, a^{(2n-1)r}\}$,
- $(b)^{SD_{8n}} \cdot (ba^2)^{SD_{8n}} = \bigcup_{r \in C_1, r \equiv 2 \pmod{4}} \{a^r, a^{(2n-1)r}\}$,
- $(ba)^{SD_{8n}} \cdot (ba)^{SD_{8n}} = \bigcup_{r \in C_1, r \equiv 2 \pmod{4}} \{a^r, a^{(2n-1)r}\}$,
- $(b)^{SD_{8n}} \cdot (ba^3)^{SD_{8n}} = \bigcup_{r \in C_2^{odd} \cup C_3^{odd}, r \equiv 3 \pmod{4}} \{a^r, a^{(2n-1)r}\}$,
- $(ba)^{SD_{8n}} \cdot (ba^2)^{SD_{8n}} = \bigcup_{r \in C_2^{odd} \cup C_3^{odd}, r \equiv 3 \pmod{4}} \{a^r, a^{(2n-1)r}\}$,

which completes the proof. \square

The Frobenius group $F_{p,q}$ can be presented by

$$F_{p,q} = \langle a, b \mid a^p = b^q = e, b^{-1}ab = a^u \rangle,$$

where $u^q \equiv 1 \pmod{p}$ [10, Definition 25.6]. Let L be the subgroup of \mathbb{Z}_p^* consisting of the powers of u and $r = (p - 1)/q$. Choose coset representatives v_1, \dots, v_r for L in \mathbb{Z}_p^* . By [10, Proposition 25.9], the

conjugacy classes of $F_{p,q}$ are as follows:

$$\begin{aligned} & \{e\}, \\ & (a^{v_i})^{F_{p,q}} = \{a^{v_i l} : l \in L\} \quad (1 \leq i \leq r), \\ & (b^n)^{F_{p,q}} = \{a^m b^n : 0 \leq m \leq p-1\} \quad (1 \leq n \leq q-1). \end{aligned}$$

Then $(b^i)^{F_{p,q}} \cdot (b^j)^{F_{p,q}} = (b^{i+j})^{F_{p,q}}$, when $i+j \neq q$. If $i+j = q$, then $(b^i)^{F_{p,q}} \cdot (b^j)^{F_{p,q}} = \bigcup_{i=1}^r (a^{v_i})^{F_{p,q}} \cup (b^{i+j})^{F_{p,q}}$. On the other hand, $(a^{v_i})^{F_{p,q}} \cdot (b^j)^{F_{p,q}} = (b^j)^{F_{p,q}}$, and $(a^{v_i})^{F_{p,q}} \cdot (a^{v_j})^{F_{p,q}} = \cup (a^{v_k})^{F_{p,q}}$ such that $v_k \equiv v_i u^m + v_j u^t \pmod{p}$, where $0 \leq m, t \leq q-1$. We have explicitly computed the set $\eta(F_{p,q})$ for several pairs of distinct primes p and q , such that $q \nmid p-1$. However, we were unable to find a general formula for $\eta(F_{p,q})$.

Question 2.8. Is it possible to find a closed formula for $\eta(F_{p,q})$?

Suppose p and q are primes, and $q|p-1$. Define:

$$S_{p,q} = \langle a, b, c \mid a^p = b^q = c^2 = e, cac = a^{-1}, bc = cb, b^{-1}ab = a^r, r^q \equiv 1 \pmod{p} \rangle.$$

Proposition 2.9. Suppose G is a group of order $2pq$, p and q , $p > q$, are odd primes, and $G \not\cong S_{p,q}$. Then

$$\eta(G) \in \left\{ \{1\}, \{1, 2, \frac{p+1}{2}\}, \{1, 2, \frac{q+1}{2}\}, \{1, 2, \frac{pq+1}{2}\}, \eta(F_{p,q}) \right\}.$$

Proof. Suppose p and q are distinct odd primes, and $p > q$. Following Zhang et al. [12], if $q \nmid p-1$, then there are four non-abelian groups of order $2pq$, and if $q|p-1$, the number of such groups is six. These groups are: $R_1 = Z_{2pq}$, $R_2 = D_{2pq}$, $R_3 = Z_q \times D_{2p}$, $R_4 = Z_p \times D_{2q}$, $R_5 = Z_2 \times F_{p,q}$, and $S_{p,q}$. The last two groups are for the case when $q|p-1$. We first notice that by Proposition 2, $\eta(R_2) = \eta(D_{2pq}) = \{1, 2, \frac{pq+1}{2}\}$. Since R_1 is abelian, $\eta(R_1) = \{1\}$. On the other hand, if G is abelian and H is an arbitrary group, then it is easy to see that $\eta(G \times H) = \eta(H)$. This implies that $\eta(R_4) = \eta(Z_p \times D_{2q}) = \eta(D_{2q}) = \{1, 2, \frac{q+1}{2}\}$, $\eta(R_3) = \eta(Z_q \times D_{2p}) = \eta(D_{2p}) = \{1, 2, \frac{p+1}{2}\}$, and $\eta(R_5) = \eta(Z_2 \times F_{p,q}) = \eta(F_{p,q})$. \square

At the end of this section, we apply [6, Theorem B] for computing $\eta(G)$, where G is a non-abelian groups of order p^3 ; p is odd. These groups can be represented by

- i. $G_1 = \langle a, b; a^{p^2} = b^p = e; b^{-1}ab = a^{1+p} \rangle$,
- ii. $G_2 = \langle a, b, z; a^p = b^p = z^p = e, az = za, bz = zb, b^{-1}ab = az \rangle$.

It is well-known that $G'_1 = Z(G_1) = \langle a^p \rangle$, and $G'_2 = Z(G_2) = \langle z \rangle$. To apply [6, Theorem B], we first compute $[x, G_1]$ and $[y, G_2]$, where $x \in G_1$ and $y \in G_2$. We have:

$$\begin{aligned} [a^i b^j, G_1] &= \{[a^i b^j, a^r b^s]; 0 \leq r \leq p^2 - 1, 0 \leq s \leq p - 1\}, \\ &= \{a^i (b^j a^r b^{-j}) (b^s a^{-i} b^{-s}) a^{-r}; 0 \leq r \leq p^2 - 1, 0 \leq s \leq p - 1\}, \\ &= \{a^i a^{r(1+p)^j} a^{-i(1+p)^s} a^{-r}; 0 \leq r \leq p^2 - 1, 0 \leq s \leq p - 1\}, \\ &= \{a^{p(rj-is)}; 0 \leq r \leq p^2 - 1, 0 \leq s \leq p - 1\} \end{aligned}$$

and

$$\begin{aligned} [a^i b^j z^k, G_2] &= \{[a^i b^j z^k, a^r b^s z^t]; 0 \leq r, s \leq p - 1\}, \\ &= \{(a^i b^j z^k) (a^r b^s z^t) (a^i b^j z^k)^{-1} (a^r b^s z^t)^{-1}; 0 \leq r, s \leq p - 1\}, \\ &= \{a^i b^j a^r (b^{s-j} a^{-i} b^{-s}) a^{-r}; 0 \leq r, s \leq p - 1\}, \\ &= \{a^i b^j a^r (a^{-i} b^{-j} z^{(s-j)i}) a^{-r}; 0 \leq r, s \leq p - 1\}, \\ &= \{z^{(p-j)(r-i)}; 0 \leq r, s \leq p - 1\}. \end{aligned}$$

Therefore, $[x, G_1] = Z(G_1)$, and $[y, G_2] = Z(G_2)$ and, by [6, Theorem A(ii)], $x^{G_1} (x^{-1})^{G_1} = [x, G_1]$ and $y^{G_2} (y^{-1})^{G_2} = [y, G_2]$. This implies that for each $u, v \in G_1$ and $u', v' \in G_2$, $|[u, G_1] \cap v^{G_1} (v^{-1})^{G_1}| = |[u, G_1] \cap [v, G_1]| = p$, and $|[u', G_2] \cap v'^{G_2} (v'^{-1})^{G_2}| = |[u', G_2] \cap [v', G_2]| = p$. If $u, v \in G_1$ and $u', v' \in G_2$ are non-identity, then by [6, Theorem B(i)], $\eta(G_1) = \eta(G_2) = \{1, p\}$.

3. CONCLUDING REMARKS

In this paper, the set $\eta(G)$ was computed for some classes of finite groups. It seems that computing $\eta(G)$ for some known group G returns to some open questions in the number theory. For example, the group $G = S_{p,q}$ in Proposition 9 has exactly $\frac{4q^2+p-1}{2q}$ conjugacy classes. These are:

$$\begin{aligned} e^G &= \{e\}, c^G = \{c, ca, ca^2, \dots, ca^{p-1}\}, \\ (b^i)^G &= \{b^i, b^i a, b^i a^2, \dots, b^i a^{p-1}\}; 1 \leq i \leq q - 1, \\ (cb^i)^G &= \{cb^i, cb^i a, cb^i a^2, \dots, cb^i a^{p-1}\}; 1 \leq i \leq q - 1, \\ (a^i)^G &= \{a^i, a^{ir}, \dots, a^{ir^{q-1}}, a^{-i}, a^{-ir}, \dots, a^{-ir^{q-1}}\}; 1 \leq i \leq \frac{p-1}{2q}. \end{aligned}$$

Since $(a^{-i}ba^i) = a^{-i-1}(ab)a^i = a^{-i-1}(ba^r)a^i = a^{-i-1}ba^{i+r} = a^{-i-2}(ab)a^{i+r} = a^{-i-2}(ba^r)a^{i+r} = a^{-i-2}ba^{i+2r} = \dots = ba^{i(1-r)}$,

$$\begin{aligned} (c^k b^j a^i)^{-1} (b^l) (c^k b^j a^i) &= a^{-i} b^{-j} c^{-k} b^l c^k b^j a^i \\ &= a^{-i} b^{-j} b^l b^j a^i \\ &= a^{-i} b^l a^i \\ &= (a^{-i} b a^i)^l \\ &= (ba^{i(1-r)})^l = (b^l a^{i(1-r^l)}). \end{aligned}$$

On the other hand, since $(a^{-i}ba^i) = a^{-i-1}(ab)a^i = a^{-i-1}(ba^r)a^i = a^{-i-1}(b)a^{i+r} = a^{-i-2}(ab)a^{i+r} = a^{-i-2}(ba^r)a^{i+r} = a^{-i-2}ba^{i+2r} = \dots = ba^{i(1-r)}$,

$$\begin{aligned} (a^i b^j c^k)^{-1} c (a^i b^j c^k) &= c^{-k} b^{-j} a^{-i} c a^i b^j c^k \\ &= c^{-k} b^{-j} c a^i a^i b^j c^k \\ &= c^{-k} b^{-j} c a^{2i} b^j c^k \\ &= c^{-k} c b^{-j} a^{2i} b^j c^k \\ &= c^{-k+1} (b^{-j} a^{2i} b^j)^k \\ &= c^{-k} (c a^m c) c^{k-1} \\ &= c^{-k} a^{-m} c^{k-1} \\ &= c^{-k+1} a^m c^{k-2} \\ &= \vdots \\ &= c^{-k+(k-1)} a^{(-1)^k 2ir^j} c^{k-k} \\ &= c a^{(-1)^k 2ir^j}. \end{aligned}$$

By a similar argument as above, we have:

$$\begin{aligned} (a^i b^j c^k)^{-1} a^l (a^i b^j c^k) &= c^{-k} b^{-j} a^{-i} a^l a^i b^j c^k \\ &= c^{-k} b^{-j} a^l b^j c^k \\ &= c^{-k} a^{r^j l} c^k \\ &= c^{-k+1} (c^{-1} a^{r^j l} c) c^{k-1} \\ &= c^{-k+1} (c^{-1} a c)^{r^j l} c^{k-1} \\ &= c^{-k+2} a^{-r^j l} c^{k-2} \\ &= \vdots \\ &= a^{(-1)^k r^j l}. \end{aligned}$$

We are now ready to compute the product of conjugacy classes in G . We first noticed that $(ca^i)(cb^j a^l) = a^{-i} c c b^j a^l = a^{-i} c^2 b^j a^l = a^{-i} b^j a^l$

$= b^j a^{-ir^j} a^l = b^j a^{-ir^j+l}$. Thus, $c^G \cdot (cb^i)^G = \{c, ca, ca^2, \dots, ca^{p-1}\} \cdot \{cb^i, cb^i a, cb^i a^2, \dots, cb^i a^{p-1}\} = \{b^i, b^i a, b^i a^2, \dots, b^i a^{p-1}\} = (b^i)^G$. But $(ca^i)(ca^j) = (a^{-i}c)ca^j = a^{-i+j}$ and so $c^G \cdot c^G = \{c, ca, ca^2, \dots, ca^{p-1}\} \cdot \{c, ca, ca^2, \dots, ca^{p-1}\} = \{e, a, \dots, a^{p-1}, a^{-1}, \dots, a^{-(p-1)}\} = \bigcup_{i=0}^{\frac{p-1}{2q}} (a^i)^G$. Again, $(ca^i)(b^j a) = c(a^i b^j)a = c(b^j a^{lr^i})a = cb^j a^{lr^i}$ and so $c^G \cdot (b^i)^G = \{c, ca, ca^2, \dots, ca^{p-1}\} \cdot \{b^i, b^i a, b^i a^2, \dots, b^i a^{p-1}\} = (cb^i)^G$.

Next, since $(cb^i a^j)(ca^l) = cb^i (a^j c)a^l = cb^i ca^{-j} a^l = c^2 b^i a^{l-j} = b^i a^{l-j}$,

$$(cb^i)^G \cdot c^G = \{b^i, b^i a, b^i a^2, \dots, b^i a^{p-1}\} = (b^i)^G.$$

To compute $(cb^i)^G \cdot (cb^j)^G$, we noticed that $(cb^i a^l)(cb^j a^m) = cb^i (ca^{-l}) b^j a^m = c^2 b^i a^{-l} b^j a^m = b^i a^{-l} b^j a^m = b^{i+j} a^{-lr^j+m}$. Thus, if $i+j \neq q$, then $(cb^i)^G \cdot (cb^j)^G = (b^{i+j})^G$. Otherwise, $(cb^i)^G \cdot (cb^j)^G = \left(\bigcup_{1 \leq i \leq \frac{p-1}{2q}} (a_i)^G \right) \cup (b^q)^G$. A similar argument shows that when $i+j \neq q$, we have $(b^i)^G \cdot (b^j)^G = (b^{i+j})^G$, otherwise $(b^i)^G \cdot (b^j)^G = \left(\bigcup_{1 \leq i \leq \frac{p-1}{2q}} (a_i)^G \right) \cup (b^q)^G$.

On the other hand, the equalities $(b^i a^j)(ca^l) = b^i (a^j c)a^l = b^i (ca^{-j})a^l = b^i (ca^{-j})a^l = b^i (ca^{-j})a^l = cb^i a^{-j+l}$ imply that:

$$\begin{aligned} (b^i)^G \cdot c^G &= \{b^i, b^i a, b^i a^2, \dots, b^i a^{p-1}\} \cdot \{c, ca, ca^2, \dots, ca^{p-1}\} \\ &= \{cb^i, cb^i a, cb^i a^2, \dots, cb^i a^{p-1}\} = (cb^i)^G. \end{aligned}$$

Other calculations were similar and were recorded as follows:

- i. $c^G \cdot (a^i)^G = \{c, ca, ca^2, \dots, ca^{p-1}\} = c^G$,
- ii. $(b^i)^G \cdot (cb^j)^G = \{cb^{i+j}, cb^{i+j} a, cb^{i+j} a^2, \dots, cb^{i+j} a^{p-1}\} = (cb^{i+j})^G$,
- iii. $(b^i)^G \cdot (a^j)^G = \{b^i, b^i a, b^i a^2, \dots, b^i a^{p-1}\} = (b^i)^G$,
- iv. $(a^i)^G \cdot (b^j)^G = \{b^j, b^j a, b^j a^2, \dots, b^j a^{p-1}\} = (b^j)^G$,
- v. $(a^i)^G \cdot (cb^j)^G = \{cb^i, cb^i a, cb^i a^2, \dots, cb^i a^{p-1}\} = (cb^i)^G$.

Again, we were unable to compute $\eta((a_i)^G \cdot (a_i)^G)$, in general. Our calculations given above and computing by the small group library of GAP [11] show that $\{1, \frac{p+2q-1}{2q}\} \subset \eta(G)$.

Question 3.1. What is $\eta(S_{p,q})$?

Using a simple calculation, one can see that $\eta(U_{6n}) = \{1, 2\}$, $\eta(D_{10}) = \{1, 2, 3\}$, $\eta(V_{48}) = \{1, 2, 3, 4\}$, $\eta(SL(2, 3) \times Z_4) = \{1, 2, 3, 4, 5\}$, and $\eta((Z_3 \times ((Z_4 \times Z_2) \times Z_2)) \times Z_2) = \{1, 2, 3, 4, 5, 6\}$. In the small group library notation of GAP [11], $SL(2, 3) \times Z_4 = \text{SmallGroup}(96, 66)$ and $(Z_3 \times ((Z_4 \times Z_2) \times Z_2)) \times Z_2 = \text{SmallGroup}(96, 13)$. Thus, it is natural to ask the following question:

Question 3.2. Is there a group G , such that $\eta(G) = \{1, 2, \dots, n\}$, where $n \geq 7$. For which values of n , we can find a group G , such that $\eta(G) = \{1, 2, \dots, n\}$?

Acknowledgment

We are indebted to the referee for his/her suggestions and helpful remarks. The research works of the authors were partially supported by the University of Kashan, under the grant No of 364988/77.

REFERENCES

1. E. Adan–Bante, Derived length and products of conjugacy classes, *Israel J. Math.* **168** (2008) 93–100.
2. E. Adan–Bante, On nilpotent groups and conjugacy classes, *Houston J. Math.* **36** (2) (2010) 345–356.
3. E. Adan–Bante and J. M. Harris, On conjugacy classes of $SL(2, q)$, *Rev. Colombiana Mat.* **46** (2012) (2) 97–111.
4. Z. Arad, D. Chillag and G. Moran, Groups with a small covering number, In: Products of Conjugacy Classes in Groups, Lecture Notes in Math. 1112, Berlin, Springer–Verlag, pp. 222–244, 1985.
5. Z. Arad and E. Fisman, An analogy between products of two conjugacy classes and products of two irreducible characters in finite groups, *Proc. Edin. Math. Soc.* **30** (1987) 7–22.
6. M. R. Darafsheh and S. Mahmood Robati, Products of conjugacy classes and products of irreducible characters in finite groups, *Turkish J. Math.* **37** (2013) (4) 607–616.
7. M. R. Darafsheh and N. S. Poursalavati, On the existence of the orthogonal basis of the symmetry classes of tensors associated with certain groups, *SUT J. Math.* **37** (1) (2001) 1–17.
8. M. Hormozi and K. Rodtes, Symmetry classes of tensors associated with the semi–dihedral groups SD_{8n} , *Colloq. Math.* **131** (2013) (1) 59–67.
9. I. M. Isaacs, Character Theory of Finite Groups, Academic Press, New York, 1976.
10. G. James and M. Liebeck, Representations and Characters of Groups, Cambridge Univ. Press, London–New York, 1993.
11. The GAP Group, *GAP – Groups, Algorithms, and Programming*, Version 4.7.5; 2014.
12. C. Zhang, J. -X. Zhou and Y. -Q. Feng, Automoprphisms of cubic Cayley graphs of order $2pq$, *Discrete Math.* **309** (2009) 2687–2695.

Maryam Jalali

Department of Pure Mathematics, Faculty of Mathematical Sciences, University of Kashan, P.O.Box 87317-51167, Kashan, I. R. Iran

Ali Reza Ashrafi

Department of Pure Mathematics, Faculty of Mathematical Sciences, University of Kashan, P.O.Box 87317-51167, Kashan, I. R. Iran

Email: ashrafi@kashanu.ac.ir

COMPUTING THE PRODUCTS OF CONJUGACY CLASSES FOR SPECIFIC FINITE GROUPS

MARYAM JALALI-RAD AND ALI REZA ASHRAFI

محاسبه حاصل ضرب کلاس‌های تزویج برخی گروه‌های متناهی

مریم جلالی راد و علی‌رضا اشرفی
کاشان - دانشگاه کاشان - دانشکده علوم ریاضی-گروه ریاضی محض

فرض کنید G یک گروه متناهی و A و B دو کلاس تزویج از آن باشند. در این صورت تعداد کلاس‌های تزویج مشمول در AB را با $\eta(AB)$ و مجموعه $\eta(AB)$ ‌هایی که A و B روی کلاس‌های تزویج G تغییر می‌کنند را با $\eta(G)$ نشان می‌دهیم. هدف این مقاله محاسبه $\eta(G)$ ‌هایی است که در آن G گروهی غیر قابل تجزیه از مرتبه $2pq$ ، p و q اعدادی اول هستند، گروهی از مرتبه p^3 ، گروهی از مرتبه $4p$ یا G یکی از گروه‌های زیر است:
 $D_{2n}, T_{4n}, U_{4n}, V_{4n}, SD_{4n}$.

کلمات کلیدی: کلاس تزویج، زیرمجموعه G - پایا، p -گروه.